

General Topology

Swayam Chube

July 7, 2023

Contents

1	Topological Spaces	2
1.1	Closed Sets and Limit Points	6
1.2	Continuous Functions	9
1.3	Product Topology	11
1.4	Metric Topology	14
1.5	Quotient Topology	18
2	Connectedness and Compactness	21
2.1	Connected Spaces	21
2.2	Compact Spaces	24
2.3	Limit Point Compactness	29
2.4	Local Compactness	31
3	Countability and Separation Axioms	34
3.1	Countability Axioms	34
3.2	Separation Axioms	37
3.3	Normal Spaces	38
3.4	Urysohn's Lemma	39
3.5	The Urysohn Metrization Theorem	43
3.6	Tietze Extension Theorem	43
4	Nets and Filters	45
4.1	Nets	45
5	The Tychonoff Theorem	47
5.1	The Tychonoff Theorem	47
5.1.1	Filters and Ultrafilters	47
5.1.2	First Proof	49
5.1.3	Second Proof	49
5.2	Stone-Čech Compactification	50
6	Metrization and Paracompactness	52
6.1	Paracompactness	52
6.1.1	Partitions of Unity	54
7	Complete Metric Spaces and Function Spaces	56
7.1	Complete Metric Spaces	56
7.2	Completion of a Metric Space	57
7.3	Compactness in Metric Spaces	58
7.4	The Stone-Weierstrass Theorem	61
8	Baire Spaces and Dimension Theory	64

Chapter 1

Topological Spaces

Definition 1.1 (Topology, Topological Space). A topology on a set X is a collection \mathcal{T} of subsets of X having the following properties:

- \emptyset and X are in \mathcal{T}
- The union of the elements of any subcollection of \mathcal{T} is in \mathcal{T}
- The intersection of the elements of any finite subcollection of \mathcal{T} is in \mathcal{T}

A set X for which a topology \mathcal{T} has been specified is called a *topological space*.

Definition 1.2 (Open Set). Let X be a topological space with associated topology \mathcal{T} . A subset U of X is said to be open if it is an element of \mathcal{T} .

This immediately implies that both \emptyset and X are open. In fact, we shall see that they are also closed. The topology \mathcal{T} of all subsets of X is called the **discrete topology** while the topology $\mathcal{T} = \{\emptyset, X\}$ is called the **indiscrete topology** or the **trivial topology**.

Definition 1.3. Let X be a set and $\mathcal{T}, \mathcal{T}'$ be two topologies defined on X . If $\mathcal{T}' \supseteq \mathcal{T}$, we say that \mathcal{T}' is *finer* than \mathcal{T} . Further, if $\mathcal{T}' \subsetneq \mathcal{T}$, then \mathcal{T}' is said to be *strictly finer* than \mathcal{T} .

Definition 1.4 (Basis). If X is a set, a *basis* for a topology on X is a collection \mathcal{B} of subsets of X (called *basis elements*) such that

- For each $x \in X$, there is at least one basis element B containing x
- If x belongs to the intersection of two basis elements B_1 and B_2 , then there is a basis element B_3 containing x such that $B_3 \subseteq B_1 \cap B_2$.

Definition 1.5 (Generated Topology). Let \mathcal{B} be a basis for a topology on X . The *topology generated by* \mathcal{B} is defined as follows: A subset U of X is said to be open in X if for each $x \in U$, there is $B \in \mathcal{B}$ such that $x \in B \subseteq U$.

Proposition 1.6. *The collection \mathcal{T} generated by a basis \mathcal{B} is indeed a topology on X .*

Proof. Obviously $\emptyset, X \in \mathcal{T}$. Suppose $\{U_\alpha\}$ is a J indexed collection of sets in \mathcal{T} . Let $U = \bigcup_{\alpha \in J} U_\alpha$. Then, for each $x \in U$, there is an $\alpha \in J$ such that $x \in U_\alpha$ and thus, there is $B \in \mathcal{B}$ such that $x \in B \subseteq U_\alpha \subseteq U$ and thus $U \in \mathcal{T}$. Let $U_1, U_2 \in \mathcal{T}$ and $x \in U_1 \cap U_2$. Then, there exist $B_1, B_2 \in \mathcal{B}$ such that $x \in B_1 \subseteq U_1$ and $x \in B_2 \subseteq U_2$ and thus, $x \in B_1 \cap B_2 \subseteq U_1 \cap U_2$. But, by definition, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2 \subseteq U_1 \cap U_2$ and consequently $U_1 \cap U_2 \in \mathcal{T}$. This finishes the proof. ■

Lemma 1.7. *Let X be a set and \mathcal{B} be a basis for a topology \mathcal{T} on X . Then \mathcal{T} equals the collection of all unions of elements of \mathcal{B} .*

Proof. Trivially note that all elements of \mathcal{B} must be in \mathcal{T} and thus, their unions too. Conversely, let $U \in \mathcal{T}$, then for all $x \in U$, there is $B_x \in \mathcal{B}$ such that $x \in B_x \subseteq U$. It is not hard to see that $U = \bigcup_{x \in U} B_x$ and we have the desired conclusion. ■

Lemma 1.8. *Let X be a topological space. Suppose \mathcal{C} is a collection of open sets of X such that for each open set U of X and each $x \in U$, there is an element C of \mathcal{C} such that $x \in C \subseteq U$. Then \mathcal{C} is a basis for the topology of X .*

Proof. We first show that \mathcal{B} is a basis. Since X is an open set, for each $x \in X$, there is $C \in \mathcal{C}$ such that $x \in C$. Let $C_1, C_2 \in \mathcal{C}$. Since both C_1 and C_2 are given to be open, so is their intersection. Thus, for each $x \in C_1 \cap C_2$, there is $C \in \mathcal{C}$ such that $x \in C \subseteq C_1 \cap C_2$. Therefore, \mathcal{B} is a basis.

Let \mathcal{T}' be the topology generated by \mathcal{C} and \mathcal{T} be the topology associated with X . Let $U \in \mathcal{T}$, then for each $x \in U$, there is $C \in \mathcal{C}$ such that $x \in C \subseteq U$, and thus $U \in \mathcal{T}'$ by definition. Conversely, let $W \in \mathcal{T}'$. Since W can be written as a union of a collection of sets in \mathcal{C} , all of which are open, W must be open too and thus $W \in \mathcal{T}$. This finishes the proof. ■

Lemma 1.9. *Let \mathcal{B} and \mathcal{B}' be bases for the topologies \mathcal{T} and \mathcal{T}' , respectively, on X . Then, the following are equivalent:*

- \mathcal{T}' is finer than \mathcal{T}
- For each $x \in X$ and each basis element $B \in \mathcal{B}$ containing x , there is a basis element $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$

Proof. Suppose \mathcal{T}' is finer than \mathcal{T} . Then $B \in \mathcal{T}$ and thus $B \in \mathcal{T}'$. As a result, there is, by definition $B' \in \mathcal{T}'$ such that $x \in B' \subseteq B$.

Conversely, let $U \in \mathcal{T}$. Since \mathcal{B} generates \mathcal{T} , for each $x \in U$, there is an element $B \in \mathcal{B}$ such that $x \in B \subseteq U$. But due to the second condition, there is an element $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$, implying that U is in the topology generated by \mathcal{B}' , that is \mathcal{T}' . This finishes the proof. ■

Definition 1.10 (Subbasis). A *subbasis* \mathcal{S} for a topology on X is a collection of subsets of X whose union equals X . The topology generated by the subbasis \mathcal{S} is defined to be the collection \mathcal{T} of all unions of finite intersections of elements of \mathcal{S} .

Proposition 1.11. *The topology generated by \mathcal{S} is indeed a topology.*

Proof. For this, it suffices to show that the set \mathcal{B} of all finite intersections of elements of \mathcal{S} forms a basis. Since the union of all elements of \mathcal{S} equals X , for each $x \in X$, there is $S \in \mathcal{S}$ such that $x \in S$ and note that S must be an element of \mathcal{B} . Finally, since the intersection of any two elements of \mathcal{B} can trivially be written as a finite intersection of elements of \mathcal{S} , it must be an element of \mathcal{B} and we are done. ■

A *simple order* is a relation C such that

1. (Comparability) For all $x \neq y$, either xCy or yCx
2. (Non-reflexivity) For all x , it is not true that xCx
3. (Transitivity) For all x, y, z such that xCy and yCz , we have xCz

Suppose X is a set with a simple order relation, $<$. Suppose a and b are elements such that $a < b$, then there are four subsets of X that are called *intervals* determined by a and b :

$$\begin{aligned}(a, b) &= \{x \mid a < x < b\} \\ (a, b] &= \{x \mid a < x \leq b\} \\ [a, b) &= \{x \mid a \leq x < b\} \\ [a, b] &= \{x \mid a \leq x \leq b\}\end{aligned}$$

Definition 1.12 (Order Topology). Let X be a set with a simple order relation and assume X has more than one element. Let \mathcal{B} be the collection of all sets of the following types:

1. All open intervals (a, b) in X
2. All intervals of the form $[a_0, b)$ where a_0 is the smallest element (if any) of X
3. All intervals of the form $(a, b_0]$ where b_0 is the largest element (if any) of X

The collection \mathcal{B} is a basis for a topology on X which is called the *order topology*.

If X has no smallest element, there are no sets of type (2) and if X has no largest element, then there are not sets of type (3).

Definition 1.13 (Product Topology). Let X and Y be topological spaces. The *product topology* on $X \times Y$ is the topology having as basis the collection \mathcal{B} of all sets of the form $U \times V$ where U and V are open sets in X and Y respectively.

Proposition 1.14. *The collection \mathcal{B} is indeed a basis.*

Proof. The first condition is trivially satisfied since $X \times Y \in \mathcal{B}$. Suppose $x \in (U_1 \times V_1) \cap (U_2 \times V_2) = (U_1 \cap U_2) \times (V_1 \cap V_2) = U_3 \times V_3$ for some open sets U_3 and V_3 in X and Y respectively. This finishes the proof. ■

It is important to note here that *every* open set in $X \times Y$ need not be of the form $U \times V$ where U is open in X and V is open in Y . For a counterexample, consider \mathbb{R}^2 equipped with the standard topology. The unit ball $x^2 + y^2 < 1$ is open in \mathbb{R}^2 but cannot be expressed in the form $U \times V$.

Proposition 1.15. *If \mathcal{B} is a basis for the topology of X and \mathcal{C} is a basis for the topology of Y , then the collection*

$$\mathcal{D} = \{B \times C \mid B \in \mathcal{B}, C \in \mathcal{C}\}$$

is a basis for the product topology on $X \times Y$.

Proof. Let W be an open set in $X \times Y$ and $(x, y) \in W$. Then, by definition, there is $B \in \mathcal{B}$ and $C \in \mathcal{C}$ such that $(x, y) \in B \times C \subseteq W$, further, since B and C are open in X and Y respectively, $B \times C$ is also open in $X \times Y$ equipped with the product topology. Therefore, we are done due to a preceeding lemma. ■

Definition 1.16. Let $\pi_1 : X \times Y \rightarrow X$ be defined by the equation $\pi_1(x, y) = x$ and let $\pi_2 : X \times Y \rightarrow Y$ be defined by the equation $\pi_2(x, y) = y$. The maps π_1 and π_2 are called the *projections* of $X \times Y$ onto its first and second factors, respectively.

Then, by definition if U is an open subset of X , then $\pi_1^{-1}(U) = U \times Y$ and similarly, if V is an open subset of Y , then $\pi_2^{-1}(V) = X \times V$.

Proposition 1.17. The collection

$$\mathcal{S} = \{\pi_1^{-1}(U) \mid U \text{ is open in } X\} \cup \{\pi_2^{-1}(V) \mid V \text{ is open in } Y\}$$

is a subbasis for the product topology on $X \times Y$.

Proof. Since $X \times Y \in \mathcal{S}$, the union of all elements of \mathcal{S} is $X \times Y$ and thus \mathcal{S} is a subbasis. Let \mathcal{B} be the basis generated by all finite intersections of \mathcal{S} . It suffices to show that $\mathcal{B} = \{U \times V \mid U \text{ is open in } X \text{ and } V \text{ is open in } Y\}$. For any U and V open in X and Y respectively, we may write $U \times V = (U \times Y) \cap (X \times V)$ and is therefore a member of \mathcal{B} . Conversely, the finite intersection of elements of \mathcal{S} is of the form $(U_1 \cap \dots \cap U_m) \times (V_1 \cap \dots \cap V_m)$, which is a product of two open sets and is an element of \mathcal{B} , which finishes the proof. ■

Definition 1.18. Let X be a topological space with topology \mathcal{T} . If Y is a subset of X , the collection

$$\mathcal{T}_Y = \{Y \cap U \mid U \in \mathcal{T}\}$$

is a topology on Y , called the *subspace topology*. With this topology, the topological space Y is called a *subspace* of X . Its open sets consist of all intersections of open sets of X with Y .

Proposition 1.19. \mathcal{T}_Y is a topology on Y .

Proof. Since $\emptyset \in \mathcal{T}$, $\emptyset = Y \cap \emptyset \in \mathcal{T}_Y$ and since $X \in \mathcal{T}$, $Y = Y \cap X \in \mathcal{T}_Y$. Further,

$$\bigcup_{\alpha \in J} (U_\alpha \cap Y) = Y \cap \bigcup_{\alpha \in J} U_\alpha \in \mathcal{T}_Y$$

And finally, $(Y \cap U_1) \cap (Y \cap U_2) = Y \cap (U_1 \cap U_2) \in \mathcal{T}_Y$. This finishes the proof. ■

Lemma 1.20. If \mathcal{B} is a basis for the topology of X and $Y \subseteq X$. Then the collection

$$\mathcal{B}_Y = \{B \cap Y \mid B \in \mathcal{B}\}$$

is a basis for the subspace topology on Y .

Proof. Let V be an open set in Y . Then, there is U in X such that $V = U \cap Y$. Since each $x \in V$ is an element of U , there is, by definition $B \in \mathcal{B}$ such that $x \in B \subseteq U$, consequently, $x \in B \cap Y \subseteq V$ and we are done due to a preceeding lemma. ■

Proposition 1.21. *Let Y be a subspace of X . If U is open in Y and Y is open in X , then U is open in X .*

Proof. Follows from the fact that $U = V \cap Y$ for some V that is open in X . ■

1.1 Closed Sets and Limit Points

Definition 1.22 (Closed Set). A subset A of a topological space X is said to be *closed* if the set $X \setminus A$ is open.

Theorem 1.23. *Let X be a topological space. Then the following conditions hold:*

1. \emptyset and X are closed
2. Arbitrary intersections of closed sets are closed
3. Finite unions of closed sets are closed

Proof. All follow from De Morgan's laws. ■

Proposition 1.24. *Let Y be a subspace of X . Then a set A is closed in Y if and only if it equals the intersection of a closed set of X with Y .*

Proof. If A is closed in Y then $Y \setminus A$ is open and thus, there is an open set B in X such that $Y \setminus A = Y \cap B$. Then,

$$A = Y \setminus (Y \cap B) = Y \cap (X \setminus B)$$

which finishes the proof. ■

Corollary 1.25. *Let Y be a subspace of X . If A is closed in Y and Y is closed in X , then A is closed in X .*

Proof. Trivial. ■

Definition 1.26 (Interior, Closure). Let X be a topological space and $A \subseteq X$. The *interior* of A is defined as the union of all open sets contained in A and the *closure* of A is defined as the intersection of all closed sets containing A . The interior of A is denoted by $\text{Int } A$ and the closure of A is denoted by \overline{A} .

Then, by definition, we have that

$$\text{Int } A \subseteq A \subseteq \overline{A}$$

Corollary 1.27. *Let X be a topological space. Then $A \subseteq X$ is closed if and only if $A = \overline{A}$.*

Proof. Trivial. ■

Theorem 1.28. Let Y be a subspace of X and A be a subset of Y . Let \bar{A} denote the closure of A in X . Then, the closure of A in Y is given by $\bar{A} \cap Y$.

Proof. Let \mathcal{F} be the collection of all closed sets in X containing A . Then, by a preceding theorem, we know that the set of all closed sets in Y containing A is given by $Y \cap \mathcal{F}$. And thus,

$$\bigcup_{C \in Y \cap \mathcal{F}} C = Y \cap \bigcup_{C \in \mathcal{F}} C = Y \cap \bar{A}$$

This finishes the proof. ■

Theorem 1.29. Let A be a subset of the topological space X .

- Then $x \in \bar{A}$ if and only if every open set U containing x intersects A
- Supposing the topology of X is given by a basis, then $x \in \bar{A}$ if and only if every basis element B containing x intersects A

Proof.

- Suppose $x \in \bar{A}$ and U be an open set containing x . Suppose for the sake of contradiction, there is an open set U in X that contains x but does not intersect A , in which case $X \setminus U$ is a closed set containing A and not containing x . By definition, since $\bar{A} \subseteq X \setminus U$, x may not be an element of \bar{A} , a contradiction. Conversely, suppose every open set U containing x intersects A and that $x \notin \bar{A}$. But then, the set $X \setminus \bar{A}$ is open and contains x but does not intersect A , a contradiction.
- Suppose $x \in \bar{A}$, then every open set containing x intersects A . Since all elements of \mathcal{B} are open, they intersect A . Conversely, since every open set U containing x has a basis subset B that contains x and therefore intersects A , U must intersect A . This finishes the proof. ■

We shall see that it is more natural to use the first statement of the above theorem as a substitute for the definition of the closure.

The statement “ U is an open set containing x ” is often shortened to “ U is a **neighborhood** of x ”.

Definition 1.30. If A is a subset of the topological space X and if $x \in X$, we say that x is a *limit point* or *cluster point* or *accumulation point* of A if every neighborhood of x intersects A in some point **other than x itself**.

For example every element of \mathbb{R} is a limit point of \mathbb{Q} .

Theorem 1.31. Let A be a subset of the topological space X and let A' be the set of all limit points of A . Then

$$\bar{A} = A \cup A'$$

Proof. If $x \in A'$, due to the preceding theorem, $x \in \bar{A}$ but since by definition, $A \subseteq \bar{A}$, we have that $A \cup A' \subseteq \bar{A}$.

Conversely let $x \in \bar{A}$. If $x \in A$, we are done. If not, then x is such that every open set containing x intersects A . But since $x \notin A$, the intersection must contain at least one point distinct from x , implying that $x \in A'$. This finishes the proof. ■

Corollary 1.32. A subset of a topological space is closed if and only if it contains all its limit points.

Proof. Follows from the fact that a subset A of a topological space is closed if and only if $A = \overline{A}$. ■

Theorem 1.33. Let X be a topological space and $A \subseteq X$. Then

$$\text{Int } A = X \setminus (\overline{X \setminus A})$$

Definition 1.34 (Hausdorff Spaces). A topological space X is called a *Hausdorff space* if for each pair x_1 and x_2 of distinct points of X , there exist neighborhoods U_1 and U_2 of x_1 and x_2 respectively that are disjoint.

Theorem 1.35. Every finite point set in a Hausdorff space X is closed.

Proof. It suffices to show this for a single point set, say $\{x_0\}$. For any $x \in X$ different from x_0 , there are open sets U and V such that $x_0 \in U$ and $x \in V$ and $U \cap V = \emptyset$. And thus, x may not be in the closure of $\{x_0\}$. This finishes the proof. ■

The condition that finite point sets be closed has been given its own name, the T_1 **axiom**. Note that there is a more standard version of T_1 -spaces,

Definition 1.36 (T_1 -space). Let X be a topological space. Then, X is said to be a T_1 -space if for any two distinct points $x, y \in X$, there is a neighborhood containing x but not y .

Theorem 1.37. A space is T_1 if and only if it satisfies the T_1 axiom.

Proof. Let X be a T_1 topological space, $\{x_0\} \subseteq X$, and $x \in X \setminus \{x_0\}$. Then, there is a neighborhood of x not containing x_0 , therefore, $x \notin \overline{\{x_0\}}$, as a result, $\{x_0\}$ is closed in X , consequently, every finite point set is closed in X .

Conversely, suppose X satisfies the T_1 axiom and $x, y \in X$ be distinct. Then, $x \notin \overline{\{y\}}$, then, using the above theorems, there is a neighborhood containing x that does not contain y , equivalently, X is a T_1 -space. ■

Corollary 1.38. Every Hausdorff space is T_1 . Note that Hausdorff spaces are also known as T_2 spaces.

The converse is not true. That is, not every Fréchet space is Hausdorff. For example, let X be an infinite set and \mathcal{T} be the *co-finite* topology on X , that is,

$$\mathcal{T} = \{U \mid U^c \text{ is finite}\}$$

That the above is a topology is trivially verified. Let $x, y \in X$, then $X \setminus \{y\}$ is an open set containing x but not y , therefore (X, \mathcal{T}) is T_1 (Fréchet).

Suppose there were disjoint open sets U and V such that $x \in U \subseteq X \setminus \{y\}$ and $y \in V \subseteq X \setminus \{x\}$. But then, $V \subseteq U^c$, contradicting the finiteness of U^c . As a result, (X, \mathcal{T}) is not T_2 (Hausdorff).

Theorem 1.39. Let X be a space satisfying the T_1 axiom and $A \subseteq X$. Then the point x is a limit point of A if and only if every neighborhood of x contains infinitely many points of A .

Proof. If every neighborhood of x intersects A at infinitely many points, then it intersects it in at least one point other than x and thus $x \in A'$.

Conversely, suppose x is a limit point of x but there is a neighborhood U of x that intersects A in only finitely many points. Let $U \cap (A \setminus \{x\}) = \{x_1, \dots, x_m\}$. Then, the open set $U \cap (X \setminus \{x_1, \dots, x_m\})$ contains x but does not intersect A , which is contradictory to the fact that x is a limit point of A . ■

Theorem 1.40. If X is a Hausdorff space, then a sequence of points of X converges to at most one point of X .

Proof. Suppose the sequence $\{x_n\}$ converges to two distinct points x and y . Then, by definition, there exist disjoint neighborhoods U and V of x and y respectively. Since x_n converges to x , U contains all but finitely many elements of the sequence but that means V cannot, a contradiction. ■

1.2 Continuous Functions

Definition 1.41 (Continuity). Let X and Y be topological spaces. A function $f : X \rightarrow Y$ is said to be continuous if for each open subset V of Y , the set $f^{-1}(V)$ is open in X .

We note here that it suffices to check the above condition for just elements of either a *basis* or a *subbasis*.

Conversely, note that it need not be the case that an open set in X is mapped to an open set in Y . Simply consider any *constant function* from $\mathbb{R} \rightarrow \mathbb{R}$.

Theorem 1.42. Let X and Y be topological spaces; let $f : X \rightarrow Y$. Then the following are equivalent

1. f is continuous
2. for every subset A of X , one has $f(\overline{A}) \subseteq \overline{f(A)}$
3. for every closed set B of Y , the set $f^{-1}(B)$ is closed in X
4. for each $x \in X$ and each neighborhood V of $f(x)$, there is a neighborhood U of x such that $f(U) \subseteq V$

Proof. (1) \Rightarrow (2). Let $x \in \overline{A}$ and V be an open set containing $f(x)$. We know by definition that $f^{-1}(V)$ is open and therefore intersects A . As a consequence, V intersects $f(A)$, implying that $f(x) \in \overline{f(A)}$.

(2) \Rightarrow (3). Let $A = f^{-1}(B)$. Let $x \in \overline{A}$. Then,

$$f(x) \in f(\overline{A}) \subseteq \overline{f(A)} \subseteq \overline{B} = B$$

and thus $x \in f^{-1}(B) = A$, implying that $A \subseteq \overline{A} \subseteq A$, finishing the proof.

(3) \Rightarrow (1). Let V be an open set in Y and let $U = f^{-1}(V)$. Since $Y \setminus V$ is closed, so is $f^{-1}(Y \setminus V) = f^{-1}(Y) \setminus U = X \setminus U$. Then, by definition, U must be open.

(1) \Leftrightarrow (4). The forward direction is trivial. Conversely, let V be an open set in Y and $U = f^{-1}(V)$. For each $x \in U$, there is an open set U_x such that $U_x \subseteq U$. Then, $U = \bigcup_{x \in U} U_x$ is open. This finishes the proof. ■

The converse of (3) is not true, consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = \frac{1}{1+x^2}$. The range of said function is $(0, 1]$, which is obviously not closed in \mathbb{R} .

Further, notice that (4) is just the topological analogue of the epsilon-delta definition of continuity.

Definition 1.43 (Homeomorphism). Let X and Y be topological spaces; let $f : X \rightarrow Y$ be a bijection. If both the function f and the inverse function $f^{-1} : Y \rightarrow X$ are continuous, then f is a *homeomorphism*.

As a result, any property of X that is entirely expressed in terms of the topology of X yields, via the correspondence f , the corresponding property for the space Y . Such a property of X is called a **topological property**.

If $f : X \rightarrow Y$ is an injective, continuous map, where X and Y are topological spaces. Let Z be the image set $f(X)$, considered as a subspace of Y ; then the function $f' : X \rightarrow Z$ obtained by restricting the range of f is bijective. If f' happens to be a homeomorphism of X with Z , we say that the map $f : X \rightarrow Y$ is a **topological imbedding** or simply an **imbedding** of X in Y .

It is important to note that a bijection $f : X \rightarrow Y$ that is continuous need not have a continuous inverse. For example, consider $f : [0, 2\pi) \rightarrow \mathbb{S}^1$, given by $f(\theta) = e^{i\theta}$. Since the unit circle is compact, but $[0, 2\pi)$ is not, the inverse may not be continuous.

Theorem 1.44. Let X, Y and Z be topological spaces

1. (Constant) If $f : X \rightarrow Y$ maps all of X to a single point of Y , then it is continuous
2. (Inclusion) If A is a subspace of X , the inclusion function $j : A \hookrightarrow X$ is continuous
3. (Composites) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then the map $g \circ f : X \rightarrow Z$ is continuous
4. (Domain Restriction) If $f : X \rightarrow Y$ is continuous, and if A is a subspace of X , then the restricted function $f|_A : A \rightarrow Y$ is continuous.
5. (Range Restriction/Expansion) Let $f : X \rightarrow Y$ be continuous. If Z is a subspace of Y containing the image set $f(X)$, then the function $g : X \rightarrow Z$ obtained by restricting the range of f is continuous. If Z is a space having Y as a subspace, then the function $h : X \rightarrow Z$ obtained by expanding the range of f is continuous.
6. (Local formulation of continuity) The map $f : X \rightarrow Y$ is continuous if X can be written as the union of open sets $\{U_\alpha\}$ such that $f|_{U_\alpha}$ is continuous for each α .

Proof.

1. Trivial
2. Trivial
3. Let V be an open set in Z . Then, $g^{-1}(V)$ is open in Y and $f^{-1} \circ g^{-1}(V)$ is open in X and thus $g \circ f$ is continuous
4. Notice that $f|_A \equiv f \circ j$
5. Let V be an open set in Z . Then, there is an open set W in Y such that $V = Z \cap W$. Since the range of f is a subset of Z , we have

$$g^{-1}(V) = g^{-1}(Z \cap W) = f^{-1}(Z \cap W) = f^{-1}(W)$$

which is open in X and thus, g is continuous. A similar argument can be applied in the second case.

6. Let V be an open set in Y , then we may write

$$f^{-1}(V) = \bigcup_{\alpha} f|_{U_{\alpha}}^{-1}(V \cap f(U_{\alpha}))$$

which is a union of a collection of open sets and is therefore open. This finishes the proof. ■

Lemma 1.45 (Pasting Lemma). Let $X = A \cup B$ where A and B are closed in X . Let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If $f(x) = g(x)$ for every $x \in A \cap B$ then f and g combine to give a continuous function $h : X \rightarrow Y$ defined as

$$h(x) = \begin{cases} f(x) & x \in A \\ g(x) & x \in B \end{cases}$$

Proof. Let C be a closed subset of Y . We then have $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$. Since f is continuous, we know that $f^{-1}(C)$ is closed in A and therefore in X similarly, so is $g^{-1}(C)$, which finishes the proof. ■

Theorem 1.46. Let $f : A \rightarrow X \times Y$ be given by the equation $f(a) = (f_1(a), f_2(a))$ then f is continuous if and only if the functions $f_1 : A \rightarrow X$ and $f_2 : A \rightarrow Y$ are continuous. The maps f_1 and f_2 are called the coordinate maps of f .

Proof. We know that the projection maps π_1, π_2 are continuous. We note that $f_1(a) = \pi_1(f(a))$ and $f_2(a) = \pi_2(f(a))$. If f is continuous, then so are f_1 and f_2 .

Conversely, suppose f_1 and f_2 are continuous and $U \times V$ be a basis element for the product topology on $X \times Y$. We know due to a preceding result that both U and V are open in X and Y respectively. Then

$$f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$$

which is an intersection of two open sets and is therefore open. ■

Example 1.47. Let X be a Hausdorff space and $A \subseteq X$ such that there is a retraction $r : X \rightarrow A$. Show that A is closed in X .

Proof. Let $x \notin A$ and $a = r(x)$. Let U and V be disjoint neighborhoods of a and x respectively. Let $W = r^{-1}(U) \cap V$. Then W contains x . Suppose $W \cap A$ is nonempty, then there is $s \in W \cap A$, therefore, $s \in r^{-1}(U)$, consequently, $s \in U$ and $s \in V$, a contradiction since $U \cap V$ is empty. Hence W is a neighborhood of x disjoint from A and A is closed. ■

Lemma 1.48. Let $p : X \rightarrow Y$ be a closed continuous surjection. Then for any $y \in Y$ and an open set U containing $p^{-1}(y)$, there is an open set W in Y containing y such that $p^{-1}(W) \subseteq U$.

Proof. Let $W = Y \setminus p(X \setminus U)$. It is not hard to argue that W satisfies the required conditions. ■

1.3 Product Topology

Definition 1.49. Let J be an index set. Given a set X , we define a J -tuple of elements of X to be a function $x : J \rightarrow X$. If α is an element of J , we often denote the value of x at α by x_α rather than $x(\alpha)$ and call it the α -th coordinate of x . We often denote the function x itself by the symbol

$$(x_\alpha)_{\alpha \in J}$$

Definition 1.50 (Cartesian Product). Let $\{A_\alpha\}_{\alpha \in J}$ be an indexed family of sets and let $X = \prod_{\alpha \in J} A_\alpha$. The cartesian product of this indexed family, denoted by

$$\prod_{\alpha \in J} A_\alpha$$

is defined to be the set of all J -tuples x of elements of X such that $x_\alpha \in A_\alpha$ for each $\alpha \in J$. That is, the set of all functions

$$x : J \rightarrow \bigcup_{\alpha \in J} A_\alpha$$

such that $x(\alpha) \in A_\alpha$ for each $\alpha \in J$.

Definition 1.51 (Box Topology). Let $\{X_\alpha\}_{\alpha \in J}$ be an indexed family of topological spaces. Let us take as a basis for a topology on the product space

$$\prod_{\alpha \in J} X_\alpha$$

the collection of all sets of the form

$$\prod_{\alpha \in J} U_\alpha$$

where U_α is open in X_α for each $\alpha \in J$. The topology generated by this basis is called the *box topology*.

Definition 1.52 (Product Topology). Let \mathcal{S}_β denote the collection

$$\mathcal{S}_\beta = \{\pi_\beta^{-1}(U_\beta) \mid U_\beta \text{ open in } X_\beta\}$$

and let \mathcal{S} denote the union of these collections

$$\mathcal{S} = \bigcup_{\beta \in J} \mathcal{S}_\beta$$

The topology generated by the subbasis \mathcal{S} is called the *product topology*. In this topology $\prod_{\alpha \in J} X_\alpha$ is called a *product space*.

It is not hard to see that \mathcal{S} is indeed a subbasis and therefore defines a topology. Let \mathcal{B} be the basis induced by \mathcal{S} . Then, any basis element is a finite intersection of elements of \mathcal{S} and eventually would have the form

$$B = \bigcap_{i=1}^n \pi_{\beta_i}^{-1}(U_{\beta_i})$$

It is then obvious that the *box topology* is finer than the *product topology* since it has more open sets. In the case of finite products of topological spaces, obviously the two of them are equal, but this is not the case for

infinite products of topological spaces, since the basis of the product topology are only finite intersections of the subbasis, implying that for any basis element of the form $B = \prod_{\alpha \in J} U_\alpha$, there exist infinitely many $\alpha \in J$ such that U_α is the entire space X_α and is therefore strictly coarser than the box topology.

As a rule of thumb:

Whenever we consider the product $\prod_{\alpha \in J} X_\alpha$, we shall assume it is given the product topology unless we specifically state otherwise.

Theorem 1.53. Suppose the topology on each space X_α is given by a basis \mathcal{B}_α . The collection of all sets of the form

$$\prod_{\alpha \in J} B_\alpha$$

where $B_\alpha \in \mathcal{B}_\alpha$ for each α will serve as a basis for the box topology on $\prod_{\alpha \in J} X_\alpha$. The collection of all sets of the same form where $B_\alpha \in \mathcal{B}_\alpha$ for finitely many indices α and $B_\alpha = X_\alpha$ for all the remaining indices will serve as a basis for the product topology $\prod_{\alpha \in J} X_\alpha$.

Proof. Straightforward. ■

Theorem 1.54. If each space X_α is a Hausdorff space, then $\prod X_\alpha$ is a Hausdorff space in both the box and product topologies.

Theorem 1.55. Let $\{X_\alpha\}$ be an indexed family of spaces and $A_\alpha \subseteq X_\alpha$ for each α . If $\prod X_\alpha$ is given by either the product or box topology, then

$$\prod \overline{A_\alpha} = \overline{\prod A_\alpha}$$

Proof. Let $x = (x_\alpha)$ be a point of $\prod \overline{A_\alpha}$ and let $U = \prod U_\alpha$ be a basis element for either the box or product topology that contains x . Since $x_\alpha \in \overline{A_\alpha}$, we know that there is $y_\alpha \in U_\alpha \cap A_\alpha$ and thus $y = (y_\alpha) \in U \cap \prod A_\alpha$.

Conversely, suppose $x = (x_\alpha) \in \overline{\prod A_\alpha}$, and let V_α be an arbitrary open set in X_α containing x_α . Since $\pi_\alpha^{-1}(V_\alpha)$ is an open set containing x , it must intersect $\prod A_\alpha$, thus, there is $y = (y_\alpha) \in \pi_\alpha^{-1}(V_\alpha) \cap \prod A_\alpha$, consequently, $y_\alpha \in V_\alpha \cap A_\alpha$, and it follows that $x_\alpha \in \overline{A_\alpha}$. This completes the proof. ■

Theorem 1.56. Let $f : A \rightarrow \prod_{\alpha \in J} X_\alpha$ be given by the equation

$$f(a) = (f_\alpha(a))_{\alpha \in J}$$

where $f_\alpha : A \rightarrow X_\alpha$ for each α . Let $\prod X_\alpha$ have the product topology. Then the function f is continuous if and only if each coordinate function f_α is continuous.

Proof. First, suppose f is continuous. Let U_β be open in X_β . The function π_β^{-1} maps U_β to an open set in $\prod X_\alpha$ and is therefore continuous. As a result, $f_\beta = \pi_\beta \circ f$ is continuous.

Conversely, suppose each coordinate function f_α is continuous. We remarked earlier that $\pi_\beta^{-1}(U_\beta)$ for some open set U_β is a subbasis for the product topology and it suffices to show that the inverse image under f of the same is open to imply continuity. Indeed,

$$f^{-1} \circ \pi_\beta^{-1}(U_\beta) = f_\beta^{-1}(U_\beta)$$

which is obviously open, since f_β is known to be continuous. This finishes the proof. ■

Caution. It is important to note that the above theorem **does not** hold for the box topology. As a simple counter example, consider the box topology on \mathbb{R}^ω and the function $f : \mathbb{R} \rightarrow \mathbb{R}^\omega$ given by $f(t) = (t, t, \dots)$. Suppose f were continuous, then the inverse image of each basis element must be open in \mathbb{R}^ω . Indeed, consider

$$B = (-1, 1) \times \left(-\frac{1}{2}, \frac{1}{2}\right) \times \dots$$

the inverse image would have to contain some open interval $(-\delta, \delta)$ in the standard topology of \mathbb{R} , that is, $(-\delta, \delta) \subseteq f^{-1}(B)$, or equivalently, $f((-\delta, \delta)) \subseteq B$, which is absurd.

1.4 Metric Topology

Definition 1.57 (Metric). A *metric* on a set X is a function $d : X \times X \rightarrow \mathbb{R}$ such that

1. $d(x, y) \geq 0$ for all $x, y \in X$; equality holds if and only if $x = y$
2. $d(x, y) = d(y, x)$ for all $x, y \in X$
3. (Triangle Inequality) $d(x, y) + d(y, z) \geq d(x, z)$ for all $x, y, z \in X$

For $\epsilon > 0$, define the set

$$B_d(x, \epsilon) = \{y \mid d(x, y) < \epsilon\}$$

Definition 1.58 (Metric Topology). If d is a metric on the set X , then the collection of all ϵ -balls $B_d(x, \epsilon)$ for $x \in X$ and $\epsilon > 0$ is a basis for a topology on X , called the *metric topology* induced by d .

Proposition 1.59. The collection of all ϵ -balls $B_d(x, \epsilon)$ for all $x \in X$ and $\epsilon > 0$ is a basis.

Proof. The first condition is trivially satisfied. Suppose $z \in B(x, \epsilon) \cap B(y, \epsilon)$. Let $r = \frac{1}{2} \min\{\epsilon - d(x, z), \epsilon - d(y, z)\}$. It is obvious, due to the triangle inequality, that $B(z, r) \subseteq B(x, \epsilon) \cap B(y, \epsilon)$. ■

Definition 1.60 (Metrisable). If X is a topological space, X is said to be *metrisable* if there exists a metric d on the set X that induces the topology of X .

A **metric space** is a metrisable space X together with a specific metric d that gives the topology of X .

Definition 1.61. Let X be a metric space with metric d . A subset A of X is said to be *bounded* if there is some number M such that $d(a_1, a_2) \leq M$ for every pair a_1, a_2 of points of A . If A is bounded and non-empty, the *diameter* of A is defined to be

$$\text{diam}(A) = \sup\{d(a_1, a_2) \mid a_1, a_2 \in A\}$$

Proposition 1.62. Every metric space is Hausdorff.

Proof. Trivial. ■

Theorem 1.63. Let X be a metric space with metric d . Define $\bar{d} : X \times X \rightarrow \mathbb{R}$ by the equation

$$\bar{d}(x, y) = \min\{d(x, y), 1\}$$

Then \bar{d} is a metric that induces the same topology as d .

Proof. We need only check the triangle inequality. This is equivalent to

$$\bar{d}(x, y) + \bar{d}(y, z) \geq \bar{d}(x, z)$$

Obviously if either one of $\bar{d}(x, y)$ or $\bar{d}(y, z)$ is greater than or equal to 1, then we are done. If not, then

$$\bar{d}(x, y) + \bar{d}(y, z) = d(x, y) + d(y, z) \geq d(x, z) \geq \bar{d}(x, z) \geq \min\{d(x, z), 1\}$$

Let \mathcal{T} be the topology on X induced by d , having basis \mathcal{B} . Let $\bar{\mathcal{B}}$ be the set of all balls induced by \bar{d} having radius strictly less than 1. Let U be an open set in \mathcal{T} and $x \in U$, then, by definition, there is $B_d(x, \epsilon)$ in \mathcal{B} such that $x \in B_d(x, \epsilon) \subseteq U$. The ball $B_{\bar{d}}(x, \frac{1}{2} \min\{\epsilon, 1\})$ is contained in $B_d(x, \epsilon)$ and also contains x . Thus, $\bar{\mathcal{B}}$ is a basis for \mathcal{T} . This finishes the proof. ■

Definition 1.64 (Euclidean, square Metric). Given $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$, we define the *Euclidean metric* on \mathbb{R}^n by the equation

$$d(x, y) = \|x - y\| = \left((x_1 - y_1)^2 + \dots + (x_n - y_n)^2 \right)^{1/2}$$

and the *square metric* ρ by the equation

$$\rho(x, y) = \max\{|x_1 - y_1|, \dots, |x_n - y_n|\}$$

Lemma 1.65. Let d and d' be two metrics on the set X ; let \mathcal{T} and \mathcal{T}' be the topologies they induce, respectively. Then \mathcal{T}' is finer than \mathcal{T} if and only if for each $x \in X$ and each $\epsilon > 0$, there exists $\delta > 0$ such that

$$B_{d'}(x, \delta) \subseteq B_d(x, \epsilon) \tag{1.1}$$

Proof. Suppose the $\epsilon - \delta$ condition holds. Let $B \in \mathcal{B}_d$ be a basis element for the topology induced by d and let x be an arbitrary element of B . Then, we can find ϵ such that $B_d(x, \epsilon) \subseteq B$ and thus, there exists δ such that $x \in B_{d'}(x, \delta) \subseteq B$. Taking the union of all such δ -balls for x , we have an open set in \mathcal{T}' which corresponds to a basis element for \mathcal{T} , and thus \mathcal{T}' is finer than \mathcal{T} .

Conversely, suppose \mathcal{T}' is finer than \mathcal{T} , then the condition is trivially satisfied. ■

Theorem 1.66. The topologies on \mathbb{R}^n induced by the Euclidean metric d and the square metric ρ are the same as the product topology on \mathbb{R}^n .

Proof. We shall first show that the topologies induced by d and ρ on \mathbb{R}^n are identical. Indeed, we have, for any two points x and y that

$$\rho(x, y) \leq d(x, y) \leq \sqrt{n} \rho(x, y)$$

this immediately implies the conclusion due to the preceding lemma.

Finally, we shall show that the topology induced by ρ is same as the product topology. Let $B = (a_1, b_1) \times \dots \times (a_n, b_n)$ be a basis element of the product topology and let $x \in B$, then for each i , there is an ϵ_i such

that $(x - \epsilon_i, x + \epsilon_i) \subseteq (a_i, b_i)$. Choosing $\epsilon = \min\{\epsilon_1, \dots, \epsilon_n\}$, we have that the topology induced by ρ is finer than the product topology. But since every basis element of the ρ -topology is inherently an element of the product topology, since it is a cartesian product of open intervals, it must be that the product topology is finer than the ρ -topology. This completes the proof. ■

Definition 1.67 (Uniform Metric). Given an index set J and given points $x = (x_\alpha)_{\alpha \in J}$ and $y = (y_\alpha)_{\alpha \in J}$ of \mathbb{R}^J , let us define a metric $\bar{\rho}$ given by

$$\bar{\rho}(x, y) = \sup\{\bar{d}(x_\alpha, y_\alpha) \mid \alpha \in J\}$$

where \bar{d} is the standard bounded metric on \mathbb{R} . This is called the *uniform metric* on \mathbb{R}^J and the topology it induces is called the *uniform topology*.

Theorem 1.68. \mathbb{R}^ω with the product topology is metrizable.

Proof. Let $\bar{d} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ be the standard bounded metric on \mathbb{R} . Define the function $D : \mathbb{R}^\omega \times \mathbb{R}^\omega \rightarrow \mathbb{R}_{\geq 0}$ by

$$D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \left\{ \frac{\bar{d}(x_i, y_i)}{i} \right\}$$

We shall show that D is a metric and induces the product topology on \mathbb{R}^ω .

That D is positive semi-definite and symmetric is evident. To verify the triangle inequality, note that

$$\frac{\bar{d}(x_i, z_i)}{i} \leq \frac{\bar{d}(x_i, y_i)}{i} + \frac{\bar{d}(y_i, z_i)}{i} \leq D(\mathbf{x}, \mathbf{y}) + D(\mathbf{y}, \mathbf{z})$$

Consequently, we have

$$D(\mathbf{x}, \mathbf{z}) = \sup \left\{ \frac{\bar{d}(x_i, z_i)}{i} \right\} \leq D(\mathbf{x}, \mathbf{y}) + D(\mathbf{y}, \mathbf{z})$$

All that remains to show is that D induces the product topology on \mathbb{R}^ω . Let $\mathbf{x} \in \mathbb{R}^\omega$ and $B \subseteq \mathbb{R}^\omega$ be a basis element in the product topology. Then, B is of the form

$$B = U_1 \times \dots \times U_n \times \mathbb{R} \times \mathbb{R} \times \dots$$

Where each U_i is an open set in \mathbb{R} under the standard topology. Since \bar{d} induces the standard topology on \mathbb{R} , there is r_i for each $1 \leq i \leq n$ such that $B_{\bar{d}}(x_i, r_i) \subseteq U_i$. Now, let

$$r = \min_{1 \leq i \leq n} \left\{ \frac{r_i}{i} \right\}$$

It is not hard to see that $B_D(\mathbf{x}, r) \subseteq B$, consequently, the topology induced by D is finer than the product topology.

Conversely, let U be an open set in the topology induced by D with $\mathbf{x} \in U$. Then, there is a basis element $B_D(\mathbf{x}, r)$ that is contained in U . Let N be the smallest positive integer such that $\frac{1}{N} < r$. We shall show that the open set

$$V = (x_1 - r, x_1 + r) \times \dots \times (x_N - r, x_N + r) \times \mathbb{R} \times \mathbb{R} \times \dots$$

is contained in $B_D(\mathbf{x}, r)$. Indeed, for any $\mathbf{y} \in V$, we have that $\bar{d}(x_i, y_i) \leq \min\{r, 1\}$ and therefore, $D(\mathbf{x}, \mathbf{y}) \leq r$, giving us the desired conclusion. ■

Theorem 1.69 ($\epsilon - \delta$ Theorem). Let $f : X \rightarrow Y$; let X and Y be metrizable with metrics d_X and d_Y respectively. Then continuity of f is equivalent to the requirement that given $x \in X$ and given $\epsilon > 0$, there exists $\delta > 0$ such that

$$d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$$

Proof. Suppose f is continuous and let $\epsilon > 0$ be given. Consider the set $f^{-1}(B_Y(f(x), \epsilon))$, which is open in X and contains the point x . Therefore, there exists a δ -ball centered at x . If y is in this δ -ball, then $f(y)$ is in the ϵ -ball centered at $f(x)$ as desired.

Conversely, suppose the $\epsilon - \delta$ condition holds and let V be open in Y and $x \in f^{-1}(V)$. But since $f(x) \in V$, there exists ϵ such that $B_Y(f(x), \epsilon)$ is contained in V , consequently, there exists δ such that $B_X(x, \delta)$ is contained in $f^{-1}(V)$ and thus $f^{-1}(V)$ is open. This finishes the proof. ■

Lemma 1.70 (Sequence Lemma). Let X be a topological space; let $A \subseteq X$. If there is a sequence of points of A converging to x , then $x \in \overline{A}$; the converse holds if X is metrizable.

Proof. Suppose there is a sequence of points of A converging to x , then each neighborhood of x contains a point of A and thus, by definition, $x \in \overline{A}$.

Conversely, suppose X is metrizable and $x \in \overline{A}$. For each $n \in \mathbb{N}$, consider $B_X(x, \frac{1}{n})$ which must contain at least one point of A , call it x_n . It is not hard to see that this sequence converges to x . This finishes the proof. ■

Note that the above lemma holds even after replacing *metrizable* by *Hausdorff*.

Theorem 1.71. Let $f : X \rightarrow Y$. If the function f is continuous, then for every $x \in X$ and every convergent sequence $x_n \rightarrow x$, the sequence $f(x_n)$ converges to $f(x)$. The converse holds if X is metrizable.

Proof. Suppose f is continuous. Let V be an open set in Y containing $f(x)$. Then, $U = f^{-1}(V)$ is an open set in X containing x . By definition, there is $N \in \mathbb{N}$ such that for all $n > N$, $x_n \in U$ and as a result, $f(x_n) \in V$. This finishes the proof.

Conversely, suppose X is metrizable, with metric d and for each convergent sequence $x_n \rightarrow x$, the sequence $f(x_n)$ converges to $f(x)$. Let A be a subset of X . We shall show that $f(\overline{A}) \subseteq \overline{f(A)}$, which would immediately imply continuity due to a preceding theorem. Let $x \in \overline{A}$, and let x_n be a point of A within the ball $B(x, \frac{1}{n})$. The sequence x_n converges to x and so does $f(x_n)$ to $f(x)$, as a result, for each open set containing $f(x)$, there is a point of $f(A)$ in it. This finishes the proof. ■

Definition 1.72 (Uniform Convergence). Let $f_n : X \rightarrow Y$ be a sequence of functions from the set X to the metric space Y . Let d be the metric for Y . We say that the sequence f_n converges uniformly to the function $f : X \rightarrow Y$ if given $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $d(f_n(x), f(x)) < \epsilon$ for all $n > N$ and all $x \in X$.

Theorem 1.73 (Uniform Limit Theorem). Let $f_n : X \rightarrow Y$ be a sequence of continuous functions from the topological space X to the metric space Y . If (f_n) converges uniformly to $f : X \rightarrow Y$, then f is continuous.

Proof. Let V be an open set in Y and $U = f^{-1}(V)$. Let $x_0 \in U$. We shall show that there is a neighborhood containing x_0 , that is contained in U . Let $y_0 = f(x_0)$ and $\epsilon > 0$ be such that $B(y_0, \epsilon) \subseteq V$. We know there exists $N \in \mathbb{N}$ such that for all $x \in X$, $d(f_n(x), f(x)) < \epsilon/3$ for all $n \geq N$. Further, there is an open set W in

X that contains x_0 such that $d(f_N(x_0), f_N(y)) < \epsilon/3$ for all $y \in W$, due to continuity of each f_n . Then, we have, for all $y \in W$

$$d(f(x), f(y)) \leq d(f(x_0), f_N(x_0)) + d(f_N(x_0), f_N(y)) + d(f_N(y), f(y)) < \epsilon$$

and thus, $f(y) \in V$. This finishes the proof. ■

It is not sufficient to replace *uniform convergence* with *pointwise convergence*. Take for example the sequence of functions $\{\cos^n x\}_{n=1}^\infty$ from $[-\frac{\pi}{2}, \frac{\pi}{2}]$ to $[0, 1]$. The limiting function is given by

$$f(x) = \begin{cases} 1 & x = 0 \\ 0 & \text{otherwise} \end{cases}$$

which is obviously not continuous.

1.5 Quotient Topology

Definition 1.74 (Quotient Map). Let X and Y be topological spaces and $p : X \rightarrow Y$ be a surjection. The map p is said to be a *quotient map* provided a subset U of Y is open in Y if and only if $p^{-1}(U)$ is open in X .

Obviously, p must be continuous. One notes that this condition is stronger than continuity, and is often called **strong continuity**.

Lemma 1.75. Let X and Y be topological spaces. Then $p : X \rightarrow Y$ is a quotient map if and only if it is surjective and for each $A \subseteq Y$, $p^{-1}(A)$ is closed in X if and only if A is closed in Y .

Proof. Suppose p is a quotient map. Then,

$$p^{-1}(Y \setminus A) = X \setminus p^{-1}(A)$$

If A were closed in Y , then $p^{-1}(A)$ is closed in X since p is continuous. On the other hand, if $p^{-1}(A)$ were closed in X , then $p^{-1}(Y \setminus A)$ would be open in X , and therefore, so would $Y \setminus A$, equivalently A is closed.

The converse is trivially evident, since it is equivalent to saying A is open in Y if and only if $p^{-1}(A)$ is open in X . ■

Definition 1.76 (Open, Closed Map). Let X and Y be topological spaces and $f : X \rightarrow Y$. Then f is said to be an *open map* if it maps open sets in X to open sets in Y and is said to be a *closed map* if it maps closed sets in X to closed sets in Y .

It immediately follows that $p : X \rightarrow Y$ is a quotient map if p is surjective, continuous and either open or closed.

We say a subset C of X is saturated with respect to the surjective map $p : X \rightarrow Y$ if it equals the complete inverse image of a subset of Y . Formally, there exists $A \subseteq Y$ such that $C = p^{-1}(A)$.

Definition 1.77 (Quotient Topology). Let X be a topological space, A a set and $p : X \rightarrow A$ be a surjective map. Then there exists exactly one topology \mathcal{T} on A relative to which p is a quotient map; it is called the *quotient topology* induced by p .

Proposition 1.78. *The above defined topology \mathcal{T} is indeed a topology and is unique.*

Proof. Let

$$\mathcal{T} = \{p(U) \mid U \in \mathcal{T}_X\}$$

That \mathcal{T} is indeed a topology follows from

$$\bigcup_{V \in \mathcal{T}} V = \bigcup_{U \in p^{-1}(B)} p(U) \in \mathcal{T} = p \left(\bigcup_{U \in p^{-1}(B)} U \right) \in \mathcal{T}$$

Let $\{V_i\}_{i=1}^n$ be a collection of open sets in \mathcal{T} , then, there is a collection $\{U_i\}_{i=1}^n$ of open sets in \mathcal{T}_X such that $U_i = p^{-1}(V_i)$. We then have

$$\bigcap_{i=1}^n V_i = \bigcap_{i=1}^n p(U_i) = p \left(\bigcap_{i=1}^n U_i \right) \in \mathcal{T}$$

We shall now show that \mathcal{T} is unique. Suppose \mathcal{T}' is another topology induced by p on A . It is obvious that $\mathcal{T} \subseteq \mathcal{T}'$. Further, for any $V \in \mathcal{T}'$, $U = p^{-1}(V) \in \mathcal{T}_X$, therefore, $V = p(U) \in \mathcal{T}$, consequently $\mathcal{T}' \subseteq \mathcal{T}$ and we have the desired conclusion. ■

On the topological space X , consider an equivalence relation \sim and let X/\sim denote the collection of equivalence classes under the aforementioned relation. Then, the map $p : X \rightarrow X/\sim$ taking a point to the equivalence class containing it is a surjection. Due to the preceding lemma, we may topologize X/\sim such that p is a quotient map. In this case, we say that X/\sim is a quotient space of X .

Theorem 1.79. *Let $p : X \rightarrow Y$ be a quotient map and $A \subseteq X$ be a saturated subset of X with respect to p . Let $q : A \rightarrow p(A)$ denote the restriction of the map p . Then*

1. *If A is either open or closed in X , then q is a quotient map*
2. *If p is either an open or a closed map then q is a quotient map.*


Proof. 1. Suppose A is open in X . Let $U \subseteq A$ be an open set that is saturated with respect to q . Then, there is $V \subseteq p(A)$ such that $U = q^{-1}(V) = p^{-1}(V)$. Where the last equality follows from the fact that A is saturated with respect to p . Now, since p is a quotient map, by definition, V must be open in Y and in the subspace topology, it must be open in $p(A)$, consequently q maps saturated open sets to saturated open sets whence it is quotient.

2. Now suppose p is an open map and similarly, let $U \subseteq A$ be saturated with respect to q , thus, there is $V \subseteq p(A)$ with $U = q^{-1}(V) = p^{-1}(V)$. Now, there is some W open in X such that $U = W \cap A$. We have that $V = p(U) = p(W \cap A) = p(W) \cap p(A)$ where the last equality follows from the fact that A is saturated with respect to p . But since p is an open map, $p(W)$ is open in Y , consequently, V is open in $p(A)$. This completes the proof. ■

Lemma 1.80. *The composition of two quotient maps is a quotient map.*

Proof. Let $p : X \rightarrow Y$ and $q : Y \rightarrow Z$ be quotient maps. Then, the composition $r = q \circ p$ is a continuous surjection. Next, let $U \subseteq X$ be a saturated open set, that is, there is some $W \subseteq Z$ such that $V = r^{-1}(W) = p^{-1} \circ q^{-1}(W)$. As a result, V is saturated with respect to p , implying that $q^{-1}(W)$ is open and since q is an open map, W is open in Z . ■

Theorem 1.81. Let $p : X \rightarrow Y$ be a quotient map. Let Z be a topological space and $g : X \rightarrow Z$ which is constant on each fibre of Y induced by p . Then g induces a map $f : Y \rightarrow Z$ such that $f \circ p = g$. The induced map f is continuous if and only if g is continuous. Further, f is a quotient map if and only if g is a quotient map.



Proof. That f exists and is well defined is trivial. Further, if f is continuous then g is continuous and if f is quotient then g is quotient, since both properties are preserved under composition.

Conversely, suppose g is continuous. Let U be an open set in Z . We would like to show that $V = f^{-1}(U)$ is open in Y , which is equivalent to showing that $p^{-1}(V)$ is open in X . But due to the commutativity of the above diagram, $p^{-1}(V) = g^{-1}(U)$ which is open since g is continuous.

Suppose now that g is quotient. Then f is surjective and due to the commutativity of the diagram. Further, $U \subseteq Z$ is open if and only if $g^{-1}(U)$ is open in X , that is, $p^{-1}(f^{-1}(U))$ is open in X , which is equivalent to $f^{-1}(U)$ open in Y . This shows that f must be a quotient map. ■

Corollary 1.82. Let $g : X \rightarrow Z$ be a continuous surjection. Define

$$X^* = \{g^{-1}(\{z\}) \mid z \in Z\}$$

and give X^* the quotient topology with respect to the map that takes every point to its respective equivalence class.

- (a) The map g induces a bijective continuous map $f : X^* \rightarrow Z$, which is a homeomorphism if and only if g is a quotient map.
- (b) If Z is Hausdorff, then so is X^*

Proof.

- (a) That f is bijective is evident. Furthermore, since g is continuous, so is f . If f is a homeomorphism, then it is trivially a quotient map and thus g is a quotient map. On the other hand, if g is a quotient map, then so is f . A quotient map which is a bijection must be a homeomorphism (this is not hard to see).
- (b) Let $u, v \in X^*$ and $a = f(u), b = f(v)$. Then, there are disjoint open sets U and V in Z separating a and b . Then $f^{-1}(U)$ and $f^{-1}(V)$ are disjoint open sets separating u and v . ■

Chapter 2

Connectedness and Compactness

2.1 Connected Spaces

Definition 2.1 (Connected Space). Let X be a topological space. A *separation* of X is a pair U and V of disjoint nonempty open subsets of X whose union is X . The space X is said to be *connected* if there does not exist a separation of X .

The above definition can be restated as follows

A space X is connected if and only if the only subsets of X that are both open and closed in X are the empty set and X itself.

It isn't hard to show the equivalence of the two statements.

Lemma 2.2. *If Y is a subspace of X , a separation of Y is a pair of disjoint nonempty sets A and B whose union is Y , neither of which contains a limit point of the other. The space Y is connected if there exists no separation of Y .*

Proof. Suppose A and B form a separation of Y , then A is both open and closed in Y , as a result, $A = \overline{A} \cap Y = \overline{A} \cap Y$, which immediately implies $\overline{A} \cap B = \emptyset$ and vice versa.

Conversely, suppose A and B are disjoint nonempty sets whose union is Y , such that $\overline{A} \cap B = A \cap \overline{B} = \emptyset$. We may then conclude that $\overline{A} \cap Y = \emptyset$ and $\overline{B} \cap Y = B$. And thus, both A and B are closed in Y and since $A = Y \setminus B$ and $B = Y \setminus A$, they are open in Y as well and are therefore a separation of Y . This finishes the proof. ■

Lemma 2.3. *If the sets C and D form a separation of X and Y is a connected subspace of X , then Y lies entirely within C or entirely within D .*

Proof. Since C and D form a separation of X , both C and D are open in X and thus, $C \cap Y$ and $D \cap Y$ are both open in Y . If both are non-empty, then we have a separation for Y , contradicting the fact that it is connected. ■

Theorem 2.4. *The union of a collection of connected subspaces of X that have a point in common is connected.*

Proof. Let $\{A_\alpha\}$ be a collection of connected subspaces of X and $p \in \bigcup_\alpha A_\alpha$. Let $Y = \bigcup_\alpha A_\alpha$. Suppose $Y = C \cup D$ is a separation. Then, due to the preceding lemma, each of the A_α must lie in either C or D , but since they have a point p in common, they must all lie in C or all in D , and as a result, either C or D must be empty, a contradiction. ■

Theorem 2.5. *Let A be a connected subspace of X . If $A \subseteq B \subseteq \overline{A}$, then B is also connected.*

Proof. Suppose $B = C \cup D$ is a separation of B . Then, without loss of generality, A lies completely in C . Then, $\overline{A} \subseteq \overline{C}$. But due to a preceding lemma, \overline{C} and D are disjoint. This implies, B is contained entirely in \overline{C} and may not intersect D , a contradiction. ■

Theorem 2.6. *The image of a connected space under a continuous map is connected.*

Proof. Let $f : X \rightarrow Y$ be a continuous map and $Z = f(X)$. Suppose $Z = A \cup B$ is a separation of Z . Then, the sets $f^{-1}(A)$ and $f^{-1}(B)$ are open in X and are non-empty, since A and B are both within the range of f , which is Z . This contradicts the fact that X is connected. ■

Theorem 2.7. *A finite cartesian product of connected spaces is connected.*

Proof. It suffices to show the statement for the Cartesian Product of two connected spaces since the result in its generality follows due to induction. Let X and Y be connected topological spaces. Let $a \times b \in X \times Y$ be a “base point”. Note that the sets $X \times b$ and $x \times Y$ are connected for all $x \in X$. Then, we have

$$X = \bigcup_{x \in X} \underbrace{(X \times b) \cup (x \times Y)}_{T_x}$$

Further, note that all the sets T_x have the point $a \times b$ in common, as a result, their union is also connected. ■

Definition 2.8 (Linear Continuum). A simply ordered set L having more than one element is called a *linear continuum* if the following hold:

1. L has the least upper bound property
2. If $x < y$, there exists z such that $x < z < y$.

Proposition 2.9. *Let X be a well-ordered set. Then $X \times [0, 1)$ in the dictionary order is a linear continuum.*

Proof. **TODO: Add in later** ■

Theorem 2.10. *If L is a linear continuum in the order topology, then L is connected and so are intervals and rays in L .*

Proof. We shall show that every convex subspace of L is connected. Let Y be a convex subspace of L that is not connected and therefore has a separation $Y = A \cup B$. Choose $a \in A$ and $b \in B$ and let $A_0 = A \cap [a, b]$ and $B_0 = B \cap [a, b]$, each of which is open and nonempty in $[a, b]$ due to the subspace topology, which is the same as the order topology. Let $c = \sup A_0$, we know this exists because of the least upper bound property. ■

Theorem 2.11. *If X is an ordered set that is connected in the order topology, then X is a linear continuum.*

Proof. Let $x, y \in X$ with $x < y$. Suppose there is no z such that $x < z < y$, then the sets $(-\infty, y)$ and (x, ∞) form a separation of X , contradicting the connectedness.

Now, we shall show that X has the least upper bound property. Let A be a bounded subset of X . Suppose A does not have a least upper bound. Let B be the set of all upper bounds of A . We shall show that B is clopen. Let $b \in B$ since A does not have a least upper bound, there is $c \in X$ such that $c < b$ and $c \in B$. Thus (c, ∞) is an open set contained in B that contains b . Next, let $x \notin B$, then there is some $a \in A$ such that $x < a$, for if not, then x would be an upper bound for A . Then, $(-\infty, x)$ is an open set disjoint from B that contains x . As a result, B is clopen, a contradiction. ■

Theorem 2.12 (Intermediate Value Theorem). *Let $f : X \rightarrow Y$ be a continuous map, where X is a connected space and Y is an ordered set in the order topology. If a and b are two points of X and if r is a point of Y lying between $f(a)$ and $f(b)$, then there exists a point c of X such that $f(c) = r$.*

Proof. Consider the sets $A = f(X) \cap (-\infty, r)$ and $B = f(X) \cap (r, \infty)$, both of which are open in $f(X)$. Suppose there is no c such that $f(c) = r$, then $f(X) = A \cup B$, both of which are non-empty because $f(a) \in A$ and $f(b) \in B$ and is therefore a separation, a contradiction to the fact that a continuous function maps connected spaces to connected spaces. ■

Theorem 2.13. *Let $\{X_\alpha\}_{\alpha \in J}$ be a collection of connected spaces. Then the product space $\prod_{\alpha \in J} X_\alpha$ is connected.*

Proof. Fix a point $\mathbf{a} = (a_\alpha)$ in X . For each finite subset K of J , define the space $X_K := \{(x_\alpha) \mid x_\alpha = a_\alpha, \alpha \notin K\}$. Then X_K is homeomorphic to $\prod_{\alpha \in K} X_\alpha$ and therefore, connected. Let Y be the union of all such X_K for K finite. We shall show that $\bar{Y} = X$, which would imply the connectedness of X . Let $\mathbf{x} = (x_\alpha) \in X$. And $U = \prod_{\alpha \in J} U_\alpha$ be a basic open set containing \mathbf{x} . Then, there are finitely many indices $\alpha_1, \dots, \alpha_n$ such that $U_{\alpha_i} \neq X_{\alpha_i}$. Let $\mathbf{y} \in Y$ be given by

$$y_\alpha = \begin{cases} a_\alpha & \alpha \notin \{\alpha_1, \dots, \alpha_n\} \\ x_\alpha & \text{otherwise} \end{cases}$$

It follows that $\mathbf{y} \in U \cap Y$, consequently, $\mathbf{x} \in \bar{Y}$. This completes the proof. ■

Definition 2.14 (Path, Path Connected). Given points x and y of the space X , a *path* in X from x to y is a continuous map $f : [a, b] \rightarrow X$ of some closed interval in the real line into X , such that $f(a) = x$ and $f(b) = y$. A space X is said to be *path connected* if every pair of points of X can be joined by a path in X .

Proposition 2.15. *A path connected space is connected.*

Proof. Trivial. ■

Example 2.16. Let \bar{S} denote the topologist's sine curve, which is the closure of

$$\left\{ x \times \sin\left(\frac{1}{x}\right) \mid 0 < x \leq 1 \right\}$$

Then \bar{S} is connected but not path connected.

Proof. Note that $\bar{S} = S \cup \{0\} \times [-1, 1]$. Since S is the continuous image of $(0, 1]$, it is connected and therefore, so is \bar{S} . Suppose \bar{S} were path connected. Then there is a continuous function $f : [0, 1] \rightarrow \bar{S}$ such that $f(0) = 0 \times 0$ and $f(1) = 1 \times \sin 1$. Since $f^{-1}(0 \times [-1, 1])$ is closed in $[0, 1]$, it has a supremum, say a , which it contains, owing to it being closed. Then, the restriction $\tilde{f} : [a, 1] \rightarrow \bar{S}$ is such that $\tilde{f}(x) \in \{0\} \times [-1, 1]$ if and only if $x = a$. We may apply a suitable linear transformation to obtain a function $g : [0, 1] \rightarrow \bar{S}$ such that $g(x) \in \{0\} \times [-1, 1]$ if and only if $x = 0$.

We may now construct continuous functions $x, y : [0, 1] \rightarrow \mathbb{R}$ such that $g = x \times y$. Now, for any $n \in \mathbb{N}$, $x(1/n) > 0$ and hence, there is $0 < u < x(1/n)$ such that $\sin(1/u) = (-1)^n$. Due to the intermediate value theorem, there is $0 < t_n < 1/n$ such that $x(t_n) = u$. By construction, $y(t_n) = (-1)^n$. But notice that $t_n \rightarrow 0$ and since y is continuous, we must have $y(t_n) \rightarrow y(0)$, a contradiction since the sequence $\{(-1)^n\}$ does not converge. ■

Example 2.17. \mathbb{R}_K is connected but not path connected.

Proof. **\mathbb{R}_K is connected:** To do this, we show that the subspaces $(-\infty, 0)$ and $(0, \infty)$ are connected, from which we can infer that $\mathbb{R} = \overline{(-\infty, 0)} \cup \overline{(0, \infty)}$ is connected.

We contend that $(-\infty, 0)$ and $(0, \infty)$ inherit the standard topology as a subspace of \mathbb{R}_K . This is obvious for $(-\infty, 0)$. The topology inherited by $(0, \infty)$ is finer than the standard topology since \mathbb{R}_K is finer than the standard topology. Let $x \in (a, b) \setminus K$ where $0 \leq a < b$. If $x > 1$, then it is trivial to see that there is a basis element (c, d) of the standard topology such that $x \in (c, d) \subseteq (a, b) \setminus K$. On the other hand, if $x < 1$, then there is some N such that $1/(N+1) < x < 1/N$ and hence, we may choose c, d such that

$$\frac{1}{N+1} < c < x < d < \frac{1}{N}$$

Thus, we would have $x \in (c, d) \subseteq (a, b) \setminus K$. Thus, the topologies are equivalent on $(0, \infty)$.

\mathbb{R}_K is not path connected: Suppose not, then there is a continuous function $f : [0, 1] \rightarrow \mathbb{R}_K$ such that $f(0) = 0$ and $f(1) = 1$. Since $[0, 1]$ is connected and compact, so is $f([0, 1])$. Since \mathbb{R}_K is strictly finer than the standard topology, a connected subspace of \mathbb{R}_K must be an interval, since the latter are the only connected sets in the standard topology. Hence, $f([0, 1])$ is a compact connected interval in \mathbb{R}_K which contains $[0, 1]$. Since $[0, 1]$ is closed in \mathbb{R}_K and is contained in a compact interval, it must be compact. This is untrue, since $[0, 1]$ is compact Hausdorff as a subspace of the standard topology and the topology it inherits as a subspace of \mathbb{R}_K is strictly finer than the former, and therefore not compact. This completes the proof. ■

2.2 Compact Spaces

Definition 2.18 (Cover). A collection \mathcal{A} of subsets of a space X is said to *cover* X or be a *covering* of X if the union of the elements of \mathcal{A} is equal to X . It is called an *open covering* of X if its elements are open subsets of X .

Definition 2.19 (Compact). A space X is said to be compact if every open covering \mathcal{A} of X contains a finite subcollection that also covers X .

This definition is extended to subspaces through the following lemma:

Lemma 2.20. *Let Y be a subspace of X . Then Y is compact if and only if every covering of Y by sets open in X contains a finite subcollection covering Y .*

Proof. Suppose Y is compact and $\{A_\alpha\}$ is a covering of Y by sets open in X . Then, the collection $\{A_\alpha \cap Y\}$ is a covering of Y by sets open in Y and therefore has a finite subcollection $\{A_{\alpha_1} \cap Y, \dots, A_{\alpha_n} \cap Y\}$ that covers Y . As a result, $\{A_{\alpha_1}, \dots, A_{\alpha_n}\}$ is a finite subcollection of open sets in X that cover Y . The converse follows similarly. ■

Theorem 2.21 (Alexander Subbase Theorem). *Let \mathcal{S} be a subbasis for the topology on X . Then, X is compact if and only if every open cover of X by elements in \mathcal{S} has a finite subcover.*

Proof. We shall show the reverse direction since the forward one is immediate. Suppose X were not compact. Then, there is an open cover \mathcal{A} of X with no finite subcover. Let Σ be the collection of all open covers of X containing \mathcal{A} with no finite subcovers. It is not hard to show that Σ forms a poset under inclusion where every chain has an upper bound. Consequently, due to Zorn's Lemma, Σ contains a maximal element, say \mathcal{U} .

We contend that $\mathcal{S} \cap \mathcal{U}$ is an open cover for X . Suppose not, then there is $x \in X$ not covered by $\mathcal{S} \cap \mathcal{U}$. Since \mathcal{U} is an open cover, there is $U \in \mathcal{U}$ containing x . Using the fact that \mathcal{S} is a subbasis, there is a finite collection $\{S_1, \dots, S_n\}$ of elements in \mathcal{S} such that $x \in S_1 \cap \dots \cap S_n \subseteq U$.

Suppose none of the S_i 's belong to \mathcal{U} , then, due to maximality, $\mathcal{U} \cup \{S_i\}$ would admit a finite subcover, $\{V_{i,1}, \dots, V_{i,n_i}\} \cup \{S_i\}$. It is not hard to see that

$$\bigcup_{i=1}^n \{V_{i,1}, \dots, V_{i,n_i}\} \cup \{U\}$$

is an open cover of X , contradicting the construction of \mathcal{U} .

Hence, there is an index $1 \leq j \leq n$ such that $S_j \in \mathcal{U}$ and thus, X is covered by $\mathcal{S} \cap \mathcal{U}$. But since $\mathcal{S} \cap \mathcal{U} \subseteq \mathcal{S}$, it has a finite subcover, contradicting the construction of \mathcal{U} . Thus, X is compact. ■

Alternate Proof Using Ultrafilters. Suppose X is not compact. Then there is an ultrafilter \mathcal{F} on X that does not converge. We contend that $\mathcal{S} \setminus \mathcal{F}$ is an open cover for X . Let $x \in X$. Then, there is a neighborhood U of x not in \mathcal{F} . Since \mathcal{S} forms a subbasis, there is a finite collection S_1, \dots, S_n in \mathcal{S} such that $x \in S_1 \cap \dots \cap S_n \subseteq U$ whence, $S_1 \cap \dots \cap S_n \notin \mathcal{F}$. Obviously, there is S_i such that $S_i \notin \mathcal{F}$, lest $S_1 \cap \dots \cap S_n \in \mathcal{F}$. Thus, $x \in S_i \in \mathcal{S} \setminus \mathcal{F}$, implying that the latter is a cover.

According to our hypothesis, there is a finite subcollection $\{S_1, \dots, S_n\} \subseteq \mathcal{S} \setminus \mathcal{F}$ that covers X , consequently,

$$\bigcap_{i=1}^n (X \setminus S_i) = \emptyset$$

which is absurd since $X \setminus S_i \in \mathcal{F}$, and \mathcal{F} has the finite intersection property. ■

Theorem 2.22. *Every closed subspace of a compact space is compact.*

Proof. Let Y be closed in a compact space X and \mathcal{A} be an open cover for Y . The collection $\mathcal{A} \cup \{X \setminus Y\}$ is an open cover for X and therefore has a finite subcover, say \mathcal{B} . In which case, $\mathcal{B} \setminus \{X \setminus Y\}$ is a finite subcover for Y , implying that it is compact. ■

Theorem 2.23. *Every compact subspace of a Hausdorff space is closed.*

Proof. Let Y be a compact subspace of a Hausdorff space X . Let $x_0 \in X \setminus Y$. Then, for each $y \in Y$, there exist disjoint open sets U_y and V_y such that $x_0 \in U_y$ and $y \in V_y$. The collection $\mathcal{A} = \{V_y \mid y \in Y\}$ forms an open cover for Y and thus, has a finite subcover, $\{V_{y_1}, \dots, V_{y_n}\}$. The corresponding open set $\bigcap_{i=1}^n U_{y_i}$ is open in X and disjoint from each V_{y_i} and thus, disjoint from Y . This implies that for each $x_0 \in X \setminus Y$, there is an open set containing it, that is contained in $X \setminus Y$. This implies that $X \setminus Y$ is open and thus Y is closed. ■

Theorem 2.24. *Every compact subspace of a metric space is closed and bounded.*

Proof. Let (X, d) be a metric space and $A \subseteq X$ be compact. That A is closed, follows from the previous theorem. If $A = \emptyset$, then it is trivially bounded. Let $a \in A$ be any point. Notice that $\mathcal{A} = \{B(a, n) \mid n \in \mathbb{N}\}$ forms an open cover of A , and therefore has a finite subcover, implying boundedness. ■

The converse of the above theorem is not true. Consider \mathbb{R} equipped with the discrete metric. That is,

$$d(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

Note that \mathbb{R} is now bounded since it is contained in $B(0, 2)$ and is trivially closed. Furthermore, $\mathcal{A} = \{B_d(r, 0.5) \mid r \in \mathbb{R}\}$ forms an open cover for \mathbb{R} with no finite subcover since each open ball $B_d(r, 0.5)$ is singleton.

Theorem 2.25. *The image of a compact space under a continuous map is compact.*

Proof. Let $f : X \rightarrow Y$ be continuous and \mathcal{A} be an open cover for $f(X)$. Then $\mathcal{B} = \{f^{-1}(A) \mid A \in \mathcal{A}\}$ is an open cover for X and therefore has a finite subcover $\{f^{-1}(A_1), \dots, f^{-1}(A_n)\}$. This immediately implies that the collection $\{A_1, \dots, A_n\}$ is a finite subcover for $f(X)$ and thus $f(X)$ is compact. ■

Theorem 2.26. *Let $f : X \rightarrow Y$ be bijective and continuous. If X is compact and Y is Hausdorff, then f is a homeomorphism.*

Proof. Due to a preceding theorem, Y must be compact. Let U be an open set in X . It suffices to show that $f(U)$ is open in Y . Since $X \setminus U$ is closed in X , due to a preceding theorem, it must be compact, as a result, $Y \setminus f(U) = f(X \setminus U)$ must be compact and thus closed (since Y is Hausdorff). Thus, $f(U)$ is open and f is a homeomorphism. ■

Lemma 2.27 (Tube Lemma). *Let Y be a compact topological space and X be any topological space. Let N be an open set in the product topology $X \times Y$ that contains the “slice” $x \times Y$ for some $x \in X$. Then, there is an open set $W \subseteq X$ such that N contains $W \times Y$.*

Proof. For each element $y \in Y$, there is a basis element $U_y \times V_y \subseteq N$ containing $x \times y$. Therefore, $\{U_y \times V_y\}_{y \in Y}$ forms an open cover for $x \times Y$ and has a finite subcover, say $U_1 \times V_1, \dots, U_n \times V_n$. Let $W = \bigcap_{i=1}^n U_i$. Then N contains $W \times Y$, which is obviously open in the product topology. ■

Theorem 2.28. *The product of finitely many compact spaces is compact.*

Proof. It suffices to show the theorem for a product of two compact spaces since the general result follows from induction.

Let X and Y be compact spaces and \mathcal{A} be an open cover for $X \times Y$. For each $x \in X$, we note that $x \times Y$ is compact and therefore, has a finite subcover, $\{A_1, \dots, A_n\} \subseteq \mathcal{A}$. Let $N_x = \bigcup_{i=1}^n A_i$. Due to the Tube Lemma, there is an open set $W_x \subseteq X$ such that $W_x \times Y$ is contained in N_x . Finally, note that $\{W_x\}_{x \in X}$ is an open cover for X and therefore has a finite subcover, say $\{W_{x_1}, \dots, W_{x_n}\}$, consequently, $\{N_{x_1}, \dots, N_{x_n}\}$ is a finite open cover for $X \times Y$. Since each N_{x_i} is a union of a finite subset of \mathcal{A} , we have that $X \times Y$ is compact. ■

Definition 2.29 (Finite Intersection). A collection \mathcal{C} of subsets of X is said to have the finite intersection property if for every finite subcollection $\{C_1, \dots, C_n\}$, the intersection $\bigcap_{i=1}^n C_i$ is nonempty.

Theorem 2.30. *Let X be a topological space. Then X is compact if and only if for every collection \mathcal{C} of closed sets in X having the finite intersection property, the intersection $\bigcap_{C \in \mathcal{C}} C$ of all the elements of \mathcal{C} is nonempty.*

Proof. Suppose X is compact and \mathcal{C} is a collection of closed sets in X having the finite intersection property. Then, the collection $\mathcal{A} = \{X \setminus C \mid C \in \mathcal{C}\}$ consists of open sets such that no finite subcollection may cover X , due to the finite intersection property. And thus, $\bigcup_{A \in \mathcal{A}} A \subsetneq X$, and equivalently, $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$.

Conversely, let \mathcal{A} be an open cover for X and $\mathcal{C} = \{X \setminus A \mid A \in \mathcal{A}\}$. It is then obvious that $\bigcap_{C \in \mathcal{C}} C$ is empty and thus, \mathcal{C} may not have the finite intersection property. As a result, there is a finite subcollection of \mathcal{A} that covers X . This finishes the proof. ■

Lemma 2.31. *Let $f : X \rightarrow Y$ where Y is compact Hausdorff. Then f is continuous if and only if the graph of f , $G_f = \{x \times f(x) \mid x \in X\}$ is closed in $X \times Y$.*

Proof. (\implies) Suppose f is continuous. Let $x \times y \notin G_f$. Then, there are disjoint neighborhoods U and V of y and $f(x)$. Now, let $\mathcal{O} = f^{-1}(V)$, which is open in X since f is continuous. Let $x' \times y' \in \mathcal{O} \times U$. Since $f(x') \in V$ and $y' \in U$, we see that $G_f \cap \mathcal{O} \times U$ is empty, thus G_f is closed. **This direction of the proof only required Y to be Hausdorff.**

(\impliedby) Suppose G_f is closed in $X \times Y$. Let $A \subseteq Y$ be closed. Then, $X \times A$ is closed in the product topology, as a result, $G_f \cap (X \times A)$ is closed in $X \times Y$. Using the compact Hausdorff-ness of Y , we know that the projection $\pi : X \times Y \rightarrow X$ is closed and therefore, $\pi(G_f \cap (X \times A))$ is closed in X . But note that

$$\pi(G_f \cap (X \times A)) = \{x \in X \mid f(x) \in A\} = f^{-1}(A)$$

and hence, f is continuous. ■

Theorem 2.32. *Let X be a simply ordered set having the least upper bound property. In the order topology, each closed interval in X is compact.*

Proof. Let $[a, b] \subseteq X$ and \mathcal{A} be an open cover for the same. We shall show that \mathcal{A} admits a finite subcover.

Claim 1. Let $x \in [a, b)$. Then, there is $y > x$ such that $[x, y]$ can be covered by at most 2 elements of \mathcal{A} .

Proof. If x has an immediate successor, y in X , that is, an element y such that (x, y) is empty, then the closed interval $[x, y]$ can be covered by at most 2 elements of \mathcal{A} . Suppose not, then x is contained in some open set $A \in \mathcal{A}$. Since A is open, it is open in the subspace topology on $[x, b]$, consequently, it contains a basis element of the form $[x, c)$. Choose some y in $[x, c)$, the existence of which is guaranteed by the fact that x has no immediate successor. Then, $[x, y]$ is covered by the single element $A \in \mathcal{A}$.

Let \mathcal{C} be the set of all $x > a$ such that the interval $[a, x]$ can be covered by finitely many elements of \mathcal{A} . Since \mathcal{C} is non-empty, we may let $c = \sup \mathcal{C}$.

Claim 2. $c \in \mathcal{C}$

Proof. Suppose not. First, note that $c \leq b$. Therefore, there is an open set $A \in \mathcal{A}$ containing c . Consequently, it contains an interval of the form $(d, c]$. Notice that (d, c) is non-empty, for if not, then $\sup \mathcal{C} \leq d < c$. Let $e \in (d, c)$. Obviously, $e \in \mathcal{C}$, therefore, the interval $[a, e]$ can be covered by finitely many elements of \mathcal{A} and since the interval $[e, c]$ is contained in A , we conclude that $[a, c] = [a, e] \cup [e, c]$ can be covered by finitely many elements of \mathcal{A} , hence, $c \in \mathcal{C}$, a contradiction.

Finally, we shall show that $c = b$. Suppose $c < b$, then there is y satisfying $c < y \leq b$ and the interval $[c, y]$ can be covered by finitely many elements of \mathcal{A} , consequently, $[a, y] = [a, c] \cup [c, y]$ can be covered by finitely many elements of \mathcal{A} , a contradiction to the definition of c . This shows that $c = b$ and completes the proof. ■

Corollary 2.33. Let I_0^2 be the ordered square, that is, $I^2 = [0, 1] \times [0, 1]$ in the order topology. Then, I_0^2 is compact.

Theorem 2.34. A subspace A of \mathbb{R}^n is compact if and only if it is closed and is bounded in the Euclidean metric d or the square metric ρ .

Proof. It suffices to use only the ρ -metric since

$$\rho(x, y) \leq d(x, y) \leq \sqrt{n}\rho(x, y)$$

Now, suppose A is compact. The collection $\{B_\rho(0, m) \mid m \in \mathbb{N}\}$ is an open cover for A and must contain a finite subcover. Let $B_\rho(0, M)$ be the largest ball in the subcover. Since all other balls are subsets of it, the set A must be too. This implies boundedness.

Conversely, suppose A is closed and bounded. Then there exists $N \in \mathbb{N}$ such that $\rho(x, y) \leq N$ for all $x, y \in A$. Equivalently, $\rho(x, 0) \leq N$ for all $x \in A$. Thus, A is a closed subset of the compact set $[-N, N]^n$ and thus is compact due to a preceding theorem. ■

Theorem 2.35 (Extreme Value Theorem). Let $f : X \rightarrow Y$ be continuous, where Y is an ordered set in the order topology. If X is compact, then there exist points c and d in X such that $f(c) \leq f(x) \leq f(d)$ for every $x \in X$.

Proof. Since f is continuous, $A = f(X)$ is compact. Suppose A does not have a maximum element. Then, the collection

$$\mathcal{A} = \{(-\infty, a) \mid a \in A\}$$

is an open cover for A and must have a finite subcover, say

$$\{(-\infty, a_1), \dots, (-\infty, a_n)\}$$

Without loss of generality, let a_n be the maximum out of all the a_i 's. Then, we note that a_n is never covered by the subcollection, a contradiction. A similar argument may be applied for the minimum element. ■

Definition 2.36. Let (X, d) be a metric space and A be a nonempty subset of X . For each $x \in X$, define the *distance from x to A* by

$$d(x, A) = \inf\{d(x, a) \mid a \in A\}$$

Proposition 2.37. The function $d(\cdot, A) : X \rightarrow \mathbb{R}$ is a continuous function.

Proof. Let $x, y \in X$. Then, for any $a \in A$, we have

$$d(x, a) \leq d(x, y) + d(y, a)$$

Taking infimum, we see

$$d(x, A) \leq d(x, y) + d(y, A) \implies d(x, A) - d(y, A) \leq d(x, y)$$

From symmetry, we see that $|d(x, A) - d(y, A)| \leq d(x, y)$ whence the continuity follows. ■

Lemma 2.38 (Lebesgue Number Lemma). Let \mathcal{A} be an open covering of the metric space (X, d) . If X is compact, there is a $\delta > 0$ such that for each subset of X having diameter less than δ , there exists an element of \mathcal{A} containing it. The number δ is called a *Lebesgue number* for the covering \mathcal{A} .

Proof. Let \mathcal{A} be an open covering of X . If X itself is an element of \mathcal{A} then any value of δ works. Suppose now that $X \notin \mathcal{A}$ and $\{A_1, \dots, A_n\}$ be a finite subcollection of elements in \mathcal{A} that cover X and $C_i = X \setminus A_i$ for all $1 \leq i \leq n$. Define the function

$$f(x) = \frac{1}{n} \sum_{i=1}^n d(x, C_i)$$

For any $x \in X$, not all of $d(x, C_i)$ may be 0, since they cannot all share a point. Thus, $f(x) > 0$. Since X is compact and f is continuous, due to the extreme value theorem, we know that f has a minimum value, say δ . We shall show that δ is a Lebesgue number for \mathcal{A} .

Let B be a subset of X having diameter less than δ . Let $x_0 \in B$; then $B \subseteq B_d(x_0, \delta)$, further, since $f(x_0) \geq \delta$, we must have an index m such that $d(x_0, C_m) \geq \delta$. Then, obviously, $B \cap C_m = \emptyset$ and consequently, $B \subseteq A_m$. ■

Definition 2.39. Let $f : (X, d_X) \rightarrow (Y, d_Y)$ be a function. f is said to be *uniformly continuous* if given $\epsilon > 0$, there is a $\delta > 0$ such that for every pair of points $x_0, x_1 \in X$,

$$d_X(x_0, x_1) < \delta \implies d_Y(f(x_0), f(x_1)) < \epsilon$$

Theorem 2.40 (Uniform Continuity Theorem). Let $f : (X, d_X) \rightarrow (Y, d_Y)$ be a continuous map such that the metric space X is compact. Then f is uniformly continuous.

Proof. Let $\epsilon > 0$ be given. Consider the collection $\mathcal{B} = \{B_Y(y, \epsilon/2) \mid y \in Y\}$ which is an open cover of Y then $\mathcal{A} = \{f^{-1}(A) \mid A \in \mathcal{B}\}$ is an open cover of X and thus has a finite subcover, $\{A_1, \dots, A_n\}$. Let δ be the Lebesgue Number of \mathcal{A} . Then for any two points $x_0, x_1 \in X$ with $d_X(x_0, x_1) < \delta$, the two point subset $\{x_0, x_1\}$ has diameter δ and is therefore contained in some A_i . As a result, $f(x_0), f(x_1) \in B_Y(y, \epsilon/2)$ for some $y \in Y$. This immediately implies that $d_Y(f(x_0), f(x_1)) < \epsilon$. ■

2.3 Limit Point Compactness

Definition 2.41 (Limit Point Compact). A space X is said to be *limit point compact* if every infinite subset of X has a limit point.

Theorem 2.42. *Compactness implies limit point compactness.*

Proof. Let A be a set with no limit points. We shall show that A is finite. We see that A must be closed, since it trivially contains all its limit points. Since each $a \in A$ is not a limit point, we may choose an open set U_a such that $U_a \cap A = \{a\}$. Then, the collection $\mathcal{U} = \{U_a \mid a \in A\}$ is an open cover for A , consequently, $\mathcal{U} \cup \{X \setminus A\}$ is an open cover for X and has a finite subcover. Since the finite subcover can have only finitely many elements of \mathcal{U} , A must be finite. ■

Definition 2.43 (Sequentially Compact). Let X be a topological space. If (x_n) is a sequence of points of X , and if

$$n_1 < n_2 < \dots$$

is an increasing sequence of positive integers, then the sequence (x_{n_i}) is called a *subsequence* of (x_n) . The space X is said to be *sequentially compact* if every sequence of points of X has a convergent subsequence.

Theorem 2.44. *Let X be a metrizable space. Then the following are equivalent*

1. X is compact
2. X is limit point compact
3. X is sequentially compact

Proof. We have already shown that (1) \implies (2). Let us first show that (2) \implies (3). Consider the set $A = \{x_n \mid n \in \mathbb{N}\}$. If A is finite, then there is some $x \in A$ such that $x_i = x$ for infinitely many indices i . This immediately gives us a convergent subsequence. If A is infinite, then there exists $x \in X$ that is a limit point of A . Then, for each $n \in \mathbb{N}$, choose $x_n \in B(x, 1/n) \cap A$. This sequence obviously converges to x and we are done.

Finally, we show that (3) \implies (1). We first show that if X is sequentially compact, then the Lebesgue number lemma holds. Suppose not. Let \mathcal{A} be an open covering of X . Then for every positive integer n , there is a set C_n of diameter less than $1/n$ that is not contained in any element of \mathcal{A} . Choose a point $x_n \in C_n$ for all positive integers n . Since X is sequentially compact, there must exist a convergent subsequence (x_{n_i}) that converges to some point $a \in A$. Since \mathcal{A} covers X , there is some $A \in \mathcal{A}$ such that $a \in A$. Choose $\epsilon > 0$ such that $B(a, \epsilon) \subseteq A$. For sufficiently large i , we have $1/n_i < \epsilon/2$ and $d(x_{n_i}, a) < \epsilon/2$, then the set C_{n_i} lies in the $\epsilon/2$ neighborhood of x_{n_i} but since x_{n_i} lies in the $\epsilon/2$ neighborhood of a , C_{n_i} lies in the ϵ neighborhood of a , and thus $C_{n_i} \subseteq B(a, \epsilon) \subseteq A$.

Next, we show that if X is sequentially compact, then for every $\epsilon > 0$, there exists a finite covering of X by open ϵ -balls. Suppose not. Let $x_1 \in X$, then $B(x_1, \epsilon)$ may not cover X and thus, there is $x_2 \in X \setminus B(x_1, \epsilon)$. Keep choosing points in X this way, that is:

$$x_{n+1} \in X \setminus \bigcup_{i=1}^n B(x_i, \epsilon)$$

The sequence (x_n) is infinite and $d(x_i, x_j) \geq \epsilon$ whenever $i \neq j$. This obviously cannot have a convergent subsequence. A contradiction.

Coming back to the original proof. Let \mathcal{A} be an open covering for X with Lebesgue number δ . Let $\epsilon = \delta/3$. Consider the finite covering of X with ϵ -balls. Each ball has a diameter of at most $2\delta/3$ and thus

is contained in some element of \mathcal{A} . The collection of all such elements of \mathcal{A} is a finite cover of X . Thus X is compact. This finishes the proof. ■

Theorem 2.45. *Let X be a compact metric space and $f : X \rightarrow X$ be a continuous map such that $d(f(x), f(y)) < d(x, y)$ for all $x, y \in X$. Then f has a unique fixed point.*

Proof. Define $A_n = f^{(n)}(X)$. Then,

$$A_{n+1} = f^{(n)}(f(X)) \subseteq f^{(n)}(X) = A_n$$

Let $A = \bigcap_{n=1}^{\infty} A_n$. Obviously, $f(A) \subseteq A$. We shall show the reverse inclusion. Choose some $x \in A$. Then, $x \in A_{n+1}$ for all $n \in \mathbb{N}$. Hence, there is some $x_n \in X$ such that $x = f^{(n+1)}(x_n)$. Consider now the sequence $y_n = f^{(n)}(x_n)$. Since every compact metric space is sequentially compact, the sequence $\{y_n\}$ has a convergent subsequence $\{y_{n_k}\}$, converging to some $a \in X$. Then, the sequence $\{x_{n_k} = f(y_{n_k})\}$ converges to $f(a)$, since f is continuous. Finally, since $y_n \in A_n$, by definition, we see that eventually the sequence $\{y_{n_k}\}$ lies completely in A_n for all $n \in \mathbb{N}$. Then, using the fact that each A_n is closed, we must have that $a \in A_n$ for all $n \in \mathbb{N}$, whence $a \in A$.

This shows that $A = f(A)$. Suppose A had more than one point. Now, since A is closed in a compact space, it is compact and hence, there are points $x, y \in A$ such that $\text{diam}(A) = d(x, y)$. From our hypothesis, there are $x_1, y_1 \in X$ such that $f(x_1) = x$ and $f(y_1) = y$. Therefore,

$$d(x, y) = d(f(x_1), f(y_1)) < d(x_1, y_1) \leq d(x, y)$$

a contradiction. Hence, A is a singleton and contains the fixed point. ■

Note that the above result does not hold for complete metric spaces. Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = \frac{1}{2}(x + \sqrt{x^2 + 1})$$

To see that this is shrinking map, invoke the mean value theorem along with the following inequality:

$$|f'(x)| = \left| \frac{1}{2} \left(1 + \frac{x}{\sqrt{x^2 + 1}} \right) \right| < 1$$

That f does not have a fixed point is obvious.

2.4 Local Compactness

Definition 2.46 (Local Compactness). A space X is said to be *locally compact* at x if there is some compact subspace C of X that contains a neighborhood of x . If X is locally compact at each of its points, X itself is said to be *locally compact*.

One notes that a compact space is automatically locally compact. Conversely, it is not necessary that a locally compact space is compact. For example, the real line \mathbb{R} with the standard topology is locally compact but not compact.

The space \mathbb{R}^ω is *not* locally compact; none of its basis elements are contained in compact subspaces, since all basis elements are of the form

$$(a_1, b_1) \times \cdots \times (a_n, b_n) \times \mathbb{R} \times \mathbb{R} \times \cdots$$

whose closure is obviously not compact.

Theorem 2.47. *Let X be a space. Then X is locally compact Hausdorff if and only if there exists a space Y satisfying the following conditions:*

1. X is a subspace of Y
2. The set $Y \setminus X$ consists of a single point
3. Y is a compact Hausdorff space

If Y and Y' are two spaces satisfying these conditions, then there is a homeomorphism of Y with Y' that equals the identity map on X .

Proof. We first show uniqueness. Let Y and Y' be two spaces satisfying these conditions. Define the function $h : Y \rightarrow Y'$ by letting h map the single point p of $Y \setminus X$ to the single point q of $Y' \setminus X$ and letting h equal the identity on X . Obviously, h is a bijection. It suffices to show that h maps open sets in Y to open sets in Y' . Let U be open in Y . If U does not contain p , it is contained in X and is open in X . Thus, $h(U) = U$ and is open in X . But since X is open in Y' , $h(U)$ is open in Y' . Now, suppose $p \in U$. Then, $C = Y \setminus U$ is closed in Y and is thus compact in Y . Since C is contained in X , it is compact in X and thus $h(C) = C$ is compact in Y' . Since Y' is Hausdorff, C is also closed in Y' and thus $h(U) = Y' \setminus C$ is open in Y' . This establishes uniqueness.

Suppose now that X is locally compact and Hausdorff. Let $Y = X \cup \{\infty\}$. The topology on Y consists of the following sets:

1. all sets U that are open in X
2. all sets of the form $Y \setminus C$ where C is a compact subspace of X

We shall first show that this forms a topology on Y . The intersection of any two sets must be in the topology. If both sets are of the form (1), then we are trivially done. If both are of the form (2), then we have $Y \setminus C_1 \cap Y \setminus C_2 = Y \setminus (C_1 \cup C_2)$ which is obviously of the form (2). Consider an intersection of the form $U \cap (Y \setminus C) = U \cap (X \setminus C)$. Since X is Hausdorff and C is compact in X , C must also be closed in X and thus $X \setminus C$ is open in X . Now, by induction it follows that finite intersections are also elements of the topology.

We now verify arbitrary unions. Obviously arbitrary unions of sets of type (1) form sets of type (1). Arbitrary unions of sets of type (2) are of the form

$$\bigcup (Y \setminus C_\alpha) = Y \setminus \bigcap C_\alpha = Y \setminus C$$

where C is some open set in X and is therefore of type (2). Finally, we need to verify the following:

$$\left(\bigcup U_\alpha \right) \cup \left(\bigcup (Y \setminus C_\beta) \right) = U \cup (Y \setminus C) = Y \setminus (C \setminus U)$$

one notes that if C is compact in X and U is open in X , then obviously $C \setminus U$ is compact in X (the proof is Straightforward). Thus this is also of type (2). And the collection is indeed a topology.

We now show that Y is compact Hausdorff. Let $x, y \in Y$. If both lie in X , then there exist disjoint open sets U, V in X that contain x and y respectively. Now suppose $x \in X$ and $y = \infty$. Consider a compact set C in X containing a neighborhood of x . Then $Y \setminus C$ contains Y and is disjoint from said neighborhood of X and thus Y is Hausdorff. Next, suppose \mathcal{A} is an open cover of Y . Then, it must contain an element of the form $Y \setminus C$ where C is compact in X , since all the open sets in Y of type (1) do not contain ∞ . Since \mathcal{A} covers Y , $\mathcal{A} \setminus \{Y \setminus C\}$ covers C and therefore has a finite subcollection that covers C . This along with $Y \setminus C$ is a finite subcover for Y and thus Y is compact.

Finally, we show that if X is a subspace of Y satisfying all the conditions, then X is locally compact Hausdorff. The fact that X is Hausdorff follows from the fact that Y is Hausdorff. Let $x \in X$. We shall show that X is locally compact at x . Since Y is Hausdorff, there exist disjoint open sets in Y containing x and ∞ respectively. The set $Y \setminus V$ is closed in Y , but since Y is compact, $Y \setminus V$ is also compact in Y and is a subset of X that contains U . This implies local compactness and finishes the proof. ■

Definition 2.48 (Compactification). A *compactification* of a space X is a compact Hausdorff space Y containing X as a subspace such that $\overline{X} = Y$. If $Y \setminus X$ is a singleton set, then Y is called the *one-point compactification* of X . Two compactifications Y_1 and Y_2 of X are said to be *equivalent* if there is a homeomorphism $h : Y_1 \rightarrow Y_2$ inducing the identity on X .

Proposition 2.49. $[0, 1]^\omega$ in the uniform topology is not locally compact.

Proof. If $[0, 1]^\omega$ were locally compact, then there would be $\varepsilon > 0$ such that $\overline{B}(\mathbf{0}, \varepsilon)$ is compact. Consider the set of points \mathbf{a}_n in $[0, 1]^\omega$ where

$$\mathbf{a}_n(m) = \begin{cases} \varepsilon & m = n \\ 0 & \text{otherwise} \end{cases}$$

Obviously, the set $\{\mathbf{a}_1, \mathbf{a}_2, \dots\}$ does not have a limit point, consequently, $\overline{B}(\mathbf{0}, \varepsilon)$ is not limit point compact, a contradiction. ■

Chapter 3

Countability and Separation Axioms

3.1 Countability Axioms

Definition 3.1 (First Countable). A topological space X is said to have a *countable basis at x* if there is a countable collection \mathcal{B} of neighborhoods of x such that each neighborhood of x contains at least one of the elements of \mathcal{B} . A space that has a countable basis at each of its points is said to satisfy the *first countability axiom* or to be *first-countable*.

Obviously, every metrizable space satisfies this axiom, since we may take

$$\mathcal{B}_x = \left\{ B\left(x, \frac{1}{n}\right) \mid n \in \mathbb{N} \right\}$$

Theorem 3.2. Let X be a first countable topological space.

1. Let A be a subset of X . If $x \in \overline{A}$, then there is a sequence of points of A converging to x .
2. Let $f : X \rightarrow Y$. If for every convergent sequence $\{x_n\}_n$ to x , the sequence $\{f(x_n)\}_n$ converges to $f(x)$, then f is continuous.

Note that the converses for both do not require the first-countable hypothesis.

Proof.

1. Let $\mathcal{B} = \{B_1, B_2, \dots\}$ be a countable basis at x . Define the sequence $\{a_n\}_{n=1}^\infty$ by picking any point $a_n \in A \cap (B_1 \cap \dots \cap B_n)$. It is not hard to see that this sequence converges to x .
2. Let $A \subseteq X$. Then, for any $x \in \overline{A}$, there is a sequence $\{a_n\}_{n=1}^\infty$ of points in A converging to x , consequently, $\{f(a_n)\}_{n=1}^\infty$ is a sequence of points in $f(A)$ converging to $f(x)$. Therefore, $f(x) \in \overline{f(A)}$, and hence, $f(\overline{A}) \subseteq \overline{f(A)}$ and f is continuous. ■

Definition 3.3 (Second Countable). If a space X has a countable basis for its topology, then X is said to satisfy the *second countability axiom* or to be *second-countable*.

From the definition of the topology generated by a basis, *second countability* implies *first countability*. Further, not every metric space is *second-countable*. To see this, consider the following lemma:

Lemma 3.4. *Let X be a topological space with a countable basis \mathcal{B} , then any discrete subspace A of X must be countable.*

Proof. The proof follows quite naturally. Since A is discrete, for each $a \in A$, there is $B_a \in \mathcal{B}$ such that $B_a \cap A = \{a\}$. Then, the function $f : A \rightarrow \mathcal{B}$ given by $f(a) = B_a$ is an injection and thus A is countable. ■

Now, consider \mathbb{R}^ω with the uniform topology. The set $A = \{0,1\}^\omega$ is uncountable and under the uniform topology, is discrete, since $\bar{\rho}(a,b) = 1$ for all $a,b \in A$ with $a \neq b$. This immediately implies that \mathbb{R}^ω under the uniform topology may not have a countable basis and cannot be second-countable.

We now show that the same isn't true for \mathbb{R}^ω equipped with the product topology. It is well known that the countable collection of all open intervals (a,b) with both $a,b \in \mathbb{Q}$ forms a basis for \mathbb{R} . Then, \mathbb{R}^ω has a **countable basis** of all open sets of the form $\prod_{n \in \mathbb{Z}^+} U_n$ where U_n is an open interval with rational end points for finitely many values of n and $U_n = \mathbb{R}$ for all others.

Theorem 3.5. *A subspace of a first-countable space is first-countable, and a countable product of first-countable spaces is first-countable. A subspace of a second-countable space is second-countable, and a countable product of a second-countable spaces is second-countable.*

Proof. The assertion about subspaces is trivially true in both cases. As for the second part note that a cross product of countable sets is countable. ■

Definition 3.6 (Lindelöf Space). A topological space X is said to be Lindelöf if every open cover has a countable subcover.

Obviously all compact spaces are Lindelöf

Definition 3.7 (Dense). A subset A of a space X is said to be *dense* in X if $\bar{A} = X$. X is said to be separable if it has a countable dense subset.

Theorem 3.8. *The following are true:*

- (a) *The continuous image of a separable space is separable*
- (b) *An open subspace of a separable space is separable*
- (c) *Let $\{X_\alpha\}_{\alpha \in J}$ be a collection of Hausdorff spaces with at least two points each. Then $\prod_{\alpha \in J} X_\alpha$ is separable if and only if each X_α is separable and J has cardinality at most that of the continuum.*

Proof. (a) Trivial.

(b) Trivial.

(c) Suppose $X = \prod_{\alpha \in J} X_\alpha$ is separable. Since projection is a continuous map, each X_α is continuous. We shall now show that the cardinality of J is at most that of the continuum. Since each X_α is Hausdorff, with at least two points, there are disjoint open sets U_α, V_α in X_α . Let D be a countable dense subset of X . Define $D_\alpha = D \cap \pi_\alpha^{-1}(U_\alpha)$. We claim that the map $\psi : J \rightarrow 2^D$ given by $\alpha \mapsto D_\alpha$ is injective. Indeed, for $\alpha \neq \beta$, consider the open sets $U = \pi_\alpha^{-1}(U_\alpha) \cap \pi_\beta^{-1}(V_\beta)$ and $V = \pi_\beta^{-1}(U_\beta)$. Note that U and V are disjoint, further, the points in $U \cap D$ belong to D_α and due to the disjointness of U and V , they are not in $D_\beta = V \cap D$. Finally, since D is countable, the cardinality of 2^D is at most the cardinality of the continuum and thus J has cardinality at most that of the continuum.

Conversely suppose J has cardinality at most that of the continuum. Then, we may treat J as a subset of $[0, 1]$. Let $D_\alpha = \{d_{\alpha 1}, d_{\alpha 2}, \dots\}$ be a countable dense subset of X_α . Let \mathcal{J} be the countable collection of open intervals with rational endpoints in the order topology on $[0, 1]$. Consider the collection of all even length tuples of the form $(I_1, \dots, I_k; n_1, \dots, n_k)$ where the I_j 's are disjoint intervals in \mathcal{J} and n_1, \dots, n_k are positive integers. Define the point $p(I_1, \dots, I_k; n_1, \dots, n_k)$ by

$$p_\alpha = \begin{cases} d_{\alpha n_i} & \alpha \in J_i \text{ for some } i \\ d_{\alpha 1} & \text{otherwise} \end{cases}$$

Obviously, the collection of such points, D , is countable. We shall show that this collection is dense. Consider a basic open set in X , which is of the form

$$B = \pi_{\alpha_1}^{-1}(U_{\alpha_1}) \cap \dots \cap \pi_{\alpha_m}^{-1}(U_{\alpha_m})$$

Note that each U_{α_i} contains a point $d_{\alpha_i n_i}$ of D_{α_i} for some $n_i \in \mathbb{N}$. Since the α_i 's are finitely many, there are disjoint intervals I_1, \dots, I_m in \mathcal{J} containing $\alpha_1, \dots, \alpha_m$ respectively. Then, the point $p(I_1, \dots, I_m; n_1, \dots, n_m)$ belongs to B and the set D is dense as desired. ■

Theorem 3.9. Suppose that X is second countable. Then

1. X is Lindelöf
2. X is separable

Proof.

1. Let $\mathcal{B} = \{B_1, B_2, \dots\}$ be a countable basis for X and \mathcal{A} be an open cover. For each $x \in X$, let A_x be an element in \mathcal{A} containing x . By definition, there must exist a basis element B_x such that $x \in B_x \subseteq A_x$. Let $\mathcal{B} = \{B_x \mid x \in X\}$. Obviously $\mathcal{B} \subseteq \mathcal{B}$ and is therefore countable. Further, for each $B \in \mathcal{B}$, there is $A(B) \in \mathcal{A}$ containing B . Therefore, $\{A(B) \mid B \in \mathcal{B}\}$ forms a countable subcover.
2. Using the Axiom of Choice, choose a set $D = \{x_n \mid x_n \in B_n\}$. For each $x \in X \setminus D$, and an open set U containing x , then there is a basis element B_j containing x that is contained in U . Therefore, $x_j \in U$. This implies $x \in \overline{D}$. ■

Theorem 3.10. Let (X, d) be a metric space. Then, the following are equivalent

1. X is second countable
2. X is Lindelöf
3. X is separable

Proof.

- $(1) \implies (2) \wedge (1) \implies (3)$ Proved above.
- $(3) \implies (1)$ Let $D = \{x_1, x_2, \dots\}$ and $Q = \{q_1, q_2, \dots\}$. We shall show that the collection $\{B(x_i, q_j)\}_{i,j \in \mathbb{N} \times \mathbb{N}}$ is a basis for the metric topology on X .

- (2) \implies (1) Let \mathcal{A}_n denote the open cover $\{B(x, \frac{1}{n})\}_{x \in X}$. Since X is Lindelöf, it has a countable sub-cover, say \mathcal{B}_n . Define $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$ which is countable. We shall show that \mathcal{B} is a basis for the metric topology on X . Let U be a neighborhood of $x \in X$. Then, there is $r \in \mathbb{R}^+$ such that $B(x, r) \subseteq U$. Choose $N \in \mathbb{N}$ such that $\frac{1}{N} < r/2$. Let B be the element of \mathcal{B}_N that contains x . Then, for any $y \in B$, $d(x, y) \leq \frac{2}{N} < r$ and $y \in U$. Consequently, $B \subseteq U$ and we are done. ■

Proposition 3.11. \mathbb{R}_ℓ is Lindelöf.

Proof. **TODO: Add in later** ■

3.2 Separation Axioms

Definition 3.12 (Regular Spaces). Suppose one-point sets are closed in X . Then X is said to be *regular* or a T_3 -space if for each pair consisting of a point x and a closed set B disjoint from X , there exist disjoint open sets containing x and B , respectively.

Definition 3.13 (Normal Spaces). Suppose one-point sets are closed in X . Then X is said to be *normal* or a T_4 -space if for each pair A, B of disjoint closed sets in X , there exist disjoint open sets containing A and B .

It is not hard to see that

$$\text{Normal} \implies \text{Regular} \implies \text{Hausdorff}$$

Theorem 3.14. Let X be a topological space such that one point sets in X are closed.

1. X is regular if and only if given a point $x \in X$ and a neighborhood U of x , there is a neighborhood V of x such that $\overline{V} \subseteq U$
2. X is normal if and only if given a closed set A and an open set U containing A , there is an open set V containing A such that $\overline{V} \subseteq U$.

Proof.

1. Suppose X is regular and $x \in U \in \mathcal{T}_X$. Since $X \setminus U$ is closed, there are disjoint open sets V and W such that $x \in V$ and $X \setminus U \subseteq W$. It is not hard to see that $\overline{V} \cap W = \emptyset$, therefore $\overline{V} \subseteq U$.
Conversely, let $x \in U$ and $A \subseteq X$ be a closed set. Then, $X \setminus A$ is open and $x \in X \setminus A$. Therefore, there is an open set V containing x such that $\overline{V} \subseteq X \setminus A$. Then, $A \subseteq X \setminus \overline{V}$ and we are done.
2. Suppose X is normal. Then, $B = X \setminus U$ is a closed set disjoint from A . Therefore, there are open sets V, W containing A and B respectively such that $A \subseteq V$ and $B \subseteq W$. It is not hard to see that $\overline{V} \cap W = \emptyset$, therefore, $\overline{V} \subseteq U$.
Conversely, let A be closed in X . Then, $B = X \setminus U$ is closed, and the sets V and $X \setminus \overline{V}$ contain A and B respectively, and are disjoint, therefore the space is normal. ■

Theorem 3.15.

1. A subspace of a Hausdorff space is Hausdorff; a product of Hausdorff spaces is Hausdorff
2. A subspace of a regular space is regular; a product of regular spaces is regular.

Proof.

1. The subspace part is trivial. Let $(X_\alpha)_\alpha$ be a collection of Hausdorff spaces. Let $\mathbf{x}, \mathbf{y} \in \prod_\alpha X_\alpha$. Since $\mathbf{x} \neq \mathbf{y}$, there is an index β such that $x_\beta \neq y_\beta$. Therefore, there disjoint are open sets U, V in X_β such that $x_\beta \in U$ and $y_\beta \in V$. As a result, $\pi_\beta^{-1}(U)$ and $\pi_\beta^{-1}(V)$ are disjoint and open in $\prod_\alpha X_\alpha$.
2. The subspace part is trivial. Let $\mathbf{x} \in \prod_\alpha X_\alpha$ where each X_α is regular and $U \subseteq \prod_\alpha X_\alpha$ be an open set containing \mathbf{x} . Let $\prod_\alpha U_\alpha$ be a basis element of $\prod_\alpha X_\alpha$ containing \mathbf{x} that is also contained in U . For each x_α , let V_α be an open set in X_α containing it such that $\overline{V_\alpha} \subseteq U_\alpha$. Note that if $U_\alpha = X_\alpha$, choose $V_\alpha = X_\alpha$ instead. As a result, $\prod_\alpha V_\alpha$ is in the product topology and its closure is contained in U . This completes the proof. ■

3.3 Normal Spaces

Theorem 3.16. Every regular space with a countable basis is normal.

Proof. Let X be a regular space with countable basis \mathcal{B} and A, B be closed sets in X . For each $x \in X$, using regularity, there is an open set U_x containing x and disjoint from B . Further, using regularity, there is a neighborhood of x , V_x such that $\overline{V_x} \subseteq U_x$. Finally, choose a basis element B_x from \mathcal{B} containing x that is contained in V .

We now have a countable cover $\{U_n\}$ for A , such that $\overline{U_i} \cap B = \emptyset$. Similarly, choose a countable open cover $\{V_n\}$ for B , such that $\overline{V_i} \cap A = \emptyset$. Let us now define

$$U'_n = U_n \setminus \bigcup_{i=1}^n \overline{V_i} \quad V'_n = V_n \setminus \bigcup_{i=1}^n \overline{U_i}$$

We shall show that U'_i and V'_j are disjoint for any i, j . Without loss of generality, suppose $i \leq j$. Suppose $x \in U'_i \cap V'_j$, therefore, $x \in U_i$ and $x \in V_j$, but using the definition of V'_j , we must have that $x \notin V'_j$, a contradiction.

Finally, define

$$U = \bigcup_{i=1}^{\infty} U'_i \quad V = \bigcup_{j=1}^{\infty} V'_j$$

These are disjoint open sets contain A and B respectively. This concludes the proof. ■

Corollary 3.17. Every regular Lindelöf space is normal.

Proof. In the same spirit as the above proof. ■

Theorem 3.18. *Every compact Hausdorff space is normal.*

Proof. Let X be a compact Hausdorff space. We shall first show that X is regular. Indeed, let $x \in X$ and $A \subseteq X$ be a closed set. Since X is compact so is A . For all $a \in A$, there are disjoint open sets U_a and V_a such that $x \in U_a$ and $a \in V_a$. Note that $\mathcal{A} = \{U_a \mid a \in A\}$ is an open cover for A and therefore has a finite subcover $\{U_{a_1}, \dots, U_{a_n}\}$. Let

$$U = \bigcap_{i=1}^n U_{a_i} \quad V = \bigcup_{i=1}^n V_{a_i}$$

which are disjoint open sets containing x and A respectively.

Suppose A and B are disjoint closed sets in X . For each $a \in A$, there are disjoint open sets U_a and V_a such that $a \in U_a$ and $B \subseteq V_a$. Note that $\mathcal{A} = \{U_a \mid a \in A\}$ is an open cover for A , and therefore, has a finite subcover $\{U_{a_1}, \dots, U_{a_n}\}$. Choose

$$U = \bigcup_{i=1}^n U_{a_i} \quad V = \bigcap_{i=1}^n V_{a_i}$$

which are disjoint open sets containing A and B respectively. ■

Theorem 3.19. *Every metrizable space is normal.*

Proof. Let (X, d) be a metric space and A, B be two disjoint closed subsets of X . For each $a \in A$, there is ε_a such that $B(a, \varepsilon_a) \cap B = \emptyset$, similarly, for each $b \in B$, there is $\varepsilon_b > 0$ such that $B(b, \varepsilon_b) \cap A = \emptyset$. Let

$$U = \bigcup_{a \in A} B(a, \varepsilon_a/2) \quad V = \bigcup_{b \in B} B(b, \varepsilon_b/2)$$

We contend that $U \cap V = \emptyset$. Suppose not, then there is $a \in A$ and $b \in B$ such that $B(a, \varepsilon_a/2) \cap B(b, \varepsilon_b/2) \neq \emptyset$. Then, $d(a, b) < (\varepsilon_a + \varepsilon_b)/2$. But by definition, we must have $d(a, b) \geq \max\{\varepsilon_a, \varepsilon_b\}$, a contradiction. This completes the proof. ■

Theorem 3.20. *Let $p : X \rightarrow Y$ be a continuous closed surjection. If X is normal, then so is Y .*

Proof. Let A, B be disjoint open sets in Y . Then $p^{-1}(A)$ and $p^{-1}(B)$ are disjoint closed sets in X , consequently, there are disjoint open sets U and V containing $p^{-1}(A)$ and $p^{-1}(B)$ respectively.

Now, for each $a \in A$, there is a neighborhood W_a of a such that $p^{-1}(a) \subseteq W_a \subseteq U$. Similarly, there is a W_b for each $b \in B$. Define $W_A = \bigcup_{a \in A} W_a$ and $W_B = \bigcup_{b \in B} W_b$.

Then, $p^{-1}(A) \subseteq p^{-1}(W_A) \subseteq U$ and $p^{-1}(B) \subseteq p^{-1}(W_B) \subseteq V$, whence $W_A \cap W_B = \emptyset$ and the conclusion follows. ■

3.4 Urysohn's Lemma

Lemma 3.21 (Urysohn). *Let X be a normal space and $A, B \subseteq X$ be two closed sets. Then there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(A) = \{0\}$ and $f(B) = \{1\}$.*

Proof. Let P be the countable set of all rational numbers in $[0, 1]$. First, define $U_1 = X \setminus B$, which is an open set containing A . Due to the normality of X , there is an open set U_0 containing A such that $\overline{U_0} \subseteq U_1$.

We shall now define an open set U_p for all $p \in P$ such that

$$p < q \implies \overline{U}_p \subseteq U_q$$

Let P_n be the set containing the first n rational numbers in some enumeration of P such that the first two enumerated rationals are 0 and 1. Let r be the $n + 1$ -st rational in the enumeration. Obviously, since P_n is finite and $0, 1 \in P_n$, there are rationals $p, q \in P$ such that

$$\begin{aligned} p &= \max\{x \in P_n \mid x < r\} \\ q &= \min\{x \in P_n \mid x > r\} \end{aligned}$$

Now, due to the induction hypothesis, $\overline{U}_p \subseteq U_q$ and therefore, using the normality of X , there is an open set U_r such that $\overline{U}_p \subseteq U_r$ and $\overline{U}_r \subseteq U_q$.

Let $s \in P_{n+1}$. If $s < p$, $\overline{U}_s \subseteq U_p \subseteq \overline{U}_p \subseteq U_r$ and if $q < s$, $\overline{U}_r \subseteq U_q \subseteq \overline{U}_q \subseteq U_s$. Therefore, the induction hypothesis holds.

Now that we have defined U_p for all $p \in P$, we shall define

$$U_p = \begin{cases} \emptyset & p < 0 \\ X & p > 1 \end{cases}$$

Now, for all $x \in X$, define the function $f : X \rightarrow [0, 1]$ as

$$f(x) = \inf\{p \mid x \in U_p\}$$

Note that since for all $p > 1$, $x \in U_p$ and the rationals are dense in the reals, $0 \leq f(x) \leq 1$. For all $a \in A$, note that $a \in U_0$, therefore $f(a) = 0$. Similarly, for all $b \in B$, note that $b \notin U_1$, as a result $b \notin U_p$ for all $p \in [0, 1]$, but $b \in U_q$ for all $q > 1$, therefore, $f(b) = \inf\{q \in \mathbb{Q} \mid q > 1\} = 1$.

All that remains is to show that f is continuous. Let $x \in X$ and $(c, d) \in [0, 1]$ be an open interval containing $f(x)$. Choose any two rational numbers p, q such that $c < p < f(x) < q < d$. Let us consider the image of the set $Y = U_q \setminus \overline{U}_p$. For all $y \in Y$, $f(y) > p$, while $f(y) < q$, therefore, $f(y) \in (c, d)$, as a result, $f(Y) \subseteq (c, d)$ and f is continuous. This completes the proof. ■

Lemma 3.22 (Jones). *Let X be a separable and normal space. Then for any set $D \subseteq X$ that is closed and discrete in X , we have $2^{|D|} \leq 2^\omega$.*

Proof. ■

Proposition 3.23. *The Sorgenfrey plane, \mathbb{R}_ℓ^2 is not normal.*

Proof 1. Notice that $\mathbb{Q} \times \mathbb{Q}$ is dense in \mathbb{R}_ℓ^2 , which is therefore separable. Consider

$$L = \{x \times (-x) \mid x \in \mathbb{R}\} \subseteq \mathbb{R}_\ell^2$$

It is not hard to argue that L is closed in \mathbb{R}_ℓ^2 and has the discrete topology as a subspace of \mathbb{R}_ℓ^2 . Due to Lemma 3.22, we see that \mathbb{R}_ℓ^2 cannot be normal. ■

Proof 2. Let $D = \mathbb{Q} \times \mathbb{Q}$ and L be as defined in the previous proof. Suppose \mathbb{R}_ℓ^2 were normal. We shall now define a function $f : \mathcal{P}(L) \rightarrow \mathcal{P}(D)$. Let $A \subseteq L$. Then A is closed in \mathbb{R}_ℓ^2 since L is discrete and closed in \mathbb{R}_ℓ^2 . If $A = \emptyset$, define $f(A) = \emptyset$. If $A = L$, then define $f(A) = D$. If $\emptyset \subsetneq A \subsetneq L$, then there are disjoint open sets U_A and V_A containing A and $L \setminus A$ respectively. Define $f(A) = D \cap U_A$.

We contend that f is an injective function. First, note that for any $\emptyset \subsetneq A \subsetneq L$, $\emptyset \subsetneq f(A) \subsetneq L$ (this is not hard to show). Now, suppose $A, B \subseteq L$ such that $f(A) = f(B)$. If $A = \emptyset$ or $A = L$ then $B = A$. Therefore, we may suppose that $\emptyset \subsetneq A, B \subsetneq L$. Without loss of generality, suppose $A \setminus B \neq \emptyset$. Then, $U_A \cap V_B \neq \emptyset$, consequently, $f(A) \setminus f(B) \neq \emptyset$, which establishes injectivity.

The contradiction is immediate from a cardinality argument. ■

Corollary 3.24. The product of two normal spaces need not be normal. Further, \mathbb{R}_ℓ^2 is regular but not normal.

Definition 3.25. If A and B are two subsets of a topological space X , and if there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(A) = \{0\}$ and $f(B) = \{1\}$, we say that A and B can be separated by a continuous function.

Definition 3.26 (Completely Regular). A space X is *completely regular* or $T_{3\frac{1}{2}}$ if one-point sets are closed in X and for each point x_0 and each closed set A not containing x_0 , there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(x_0) = 1$ and $f(A) = \{0\}$.

Theorem 3.27. A subspace of a completely regular space is regular. A product of completely regular spaces is completely regular.

Proof. Let X be completely regular and $Y \subseteq X$. Obviously, one point sets are closed in Y . Let $x_0 \in Y$ and A be a closed set in Y that is disjoint from it. There is a closed set F in X such that $A = F \cap Y$. Then, there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(x_0) = 1$ and $f(F) = \{0\}$. Then, the restriction of f to Y is the desired continuous function separating x_0 and A .

Suppose $\{X_\alpha\}_{\alpha \in J}$ is a collection of completely regular spaces and let $X = \prod_{\alpha \in J} X_\alpha$. Let $\mathbf{a} = (a_\alpha)_{\alpha \in J}$ and A a closed set in X not containing \mathbf{a} . Then, there is a basis element $\prod_{\alpha \in J} U_\alpha$ containing \mathbf{a} that is disjoint from A . Then, there is a finite set of indices $\{\alpha_1, \dots, \alpha_n\}$ such that $U_{\alpha_i} \neq X_{\alpha_i}$ for $1 \leq i \leq n$. Since each X_{α_i} is completely regular, there is a continuous function $f_{\alpha_i} : X_{\alpha_i} \rightarrow [0, 1]$ such that $f_{\alpha_i}(a_{\alpha_i}) = 1$ and $f_{\alpha_i}(X_{\alpha_i} \setminus U_{\alpha_i}) = \{0\}$. Define the continuous function $\Phi : X \rightarrow [0, 1]$ by

$$f(\mathbf{x}) = f_{\alpha_1}(\pi_{\alpha_1}(\mathbf{x})) \cdots f_{\alpha_n}(\pi_{\alpha_n}(\mathbf{x}))$$

It is not hard to see that this function separates \mathbf{a} and A . ■

Corollary 3.28. A locally compact Hausdorff space is completely regular.

Proposition 3.29. Let X be completely regular and K, A be a disjoint compact-closed pair in X . Then, K and A can be separated by a continuous function.

Proof. For each $x \in K$, there is a continuous function $f_x : X \rightarrow [0, 1]$ such that $f(x) = 1$ and $f(A) = \{0\}$. Let $U_x := f_x^{-1}((1/2, 1])$. Then $\{U_x\}_{x \in K}$ forms an open cover for K , therefore has a finite subcover, $\{U_{x_1}, \dots, U_{x_n}\}$. Define $g = \max\{f_{x_1}, \dots, f_{x_n}\}$. Then, $g(x) > 1/2$ for all $x \in K$ and $g(A) = \{0\}$. Finally, define $h = 2 \min\{g, 1/2\}$. Then, $h(K) = \{1\}$ and $h(A) = \{0\}$. ■

Corollary 3.30. Let X be locally compact Hausdorff, K a compact subset and A a disjoint closed subset. Then, there is a continuous function separating K and A .

Theorem 3.31. *Let X be a locally compact Hausdorff space and $K \subseteq X$ be compact. Let V be an open set containing K . Then there is a continuous function $f : X \rightarrow [0, 1]$ with compact support such that $f(K) = 1$ and $\text{supp}(f) \subseteq V$.*

Proof. First, we show that there is a compact set K' and an open set U such that $K \subseteq U \subseteq K' \subseteq V$. For each $a \in K$, there is a neighborhood U_a with compact closure such that $a \in \overline{U_a} \subseteq V$. Since $\{U_a\}$ forms an open cover for K , it has a finite cover, say $\{U_{a_1}, \dots, U_{a_n}\}$. Define $U = U_{a_1} \cup \dots \cup U_{a_n}$ and $K' = \overline{U_{a_1}} \cup \dots \cup \overline{U_{a_n}}$.

Due to the previous proposition, there is a continuous function such that $f(K) = 1$ and $f(X \setminus U) = 0$. Obviously, $\{x \in X \mid f(x) \neq 0\} \subseteq K'$ and since K' is closed, $\text{Supp } f \subseteq K'$ whence it is compact. This completes the proof. ■

Lemma 3.32. *Let X be normal and $A \subseteq X$. Then, there is a function $f : X \rightarrow [0, 1]$ such that $f^{-1}(\{0\}) = A$ if and only if A is a closed G_δ set.*

Proof. Suppose there is a function $f : X \rightarrow [0, 1]$ such that $f^{-1}(\{0\}) = A$. Then, obviously A is closed, further

$$A = \bigcap_{n=1}^{\infty} f^{-1}\left(\left[0, \frac{1}{n}\right)\right)$$

whence A is G_δ .

Conversely, suppose A is a closed G_δ subset of X . Then there is a countable collection of open sets $\{U_n\}$ such that $A = \bigcap_{n=1}^{\infty} U_n$. Using normality and the Urysohn Lemma, there is a continuous function $f_n : X \rightarrow [0, 1]$ such that $f_n(A) = \{0\}$ and $f_n(X \setminus U_n) = \{1\}$. Define the function

$$f = \sum_{n=1}^{\infty} 2^{-n} f_n$$

Using the Weierstrass M-test, it is not hard to see that the convergence of the series to f is uniform and thus f is a continuous function satisfying the required properties. ■

Theorem 3.33.

Theorem 3.34 (Imbedding Theorem). *Let X be Fréchet. Suppose $\{f_\alpha\}_{\alpha \in J}$ is an indexed family of continuous functions $f_\alpha : X \rightarrow \mathbb{R}$ satisfying the requirement that for each point $x_0 \in X$ and each neighborhood U of x_0 , there is an index α such that f_α is positive at x_0 and vanishes outside U . Then the function $F : X \rightarrow \mathbb{R}^J$ defined by*

$$F(x) = (f_\alpha(x))_{\alpha \in J}$$

is an imbedding of X in \mathbb{R}^J . In particular, if f_α maps X into $[0, 1]$ for each α , then F imbeds X in $[0, 1]^J$.

Proof. That F is a continuous function is obvious. Let $Z = F(X)$. Let U be open in X . We shall show that $F(U)$ is open in Z . Choose some $z_0 \in F(U)$, then, there is some $x_0 \in U$ such that $F(x_0) = z_0$. There is some index β such that $f_\beta(x_0) > 0$ and f_β vanishes outside U . Let $W = \pi_\beta^{-1}((0, \infty)) \cap Z$. We contend that $z_0 \in W \subseteq F(U)$. That $z_0 \in W$ is obvious. Now, let $z \in W$, then there is some $x \in X$ such that $F(x) = z$. Since $\pi_\beta(z) > 0$, $f_\beta(x) > 0$ and thus $x \in U$, consequently, $F(x) \in F(U)$. This completes the proof. ■

Corollary 3.35. A space X is completely regular if and only if it is homeomorphic to a subspace of $[0, 1]^J$ for some indexing set J .

Proof. Suppose X were completely regular. For each pair (x, U) of a point and an open set U containing x , there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(x) = 1$ and $f(X \setminus U) = \{0\}$. Let $\{f_\alpha\}_{\alpha \in J}$ be the collection of all such continuous functions. Then, due to Theorem 3.34, there is an imbedding $F : X \rightarrow [0, 1]^J$. The converse is obvious. ■

Proposition 3.36. A subspace of a normal space need not be normal.

The following proof uses the forthcoming Tychonoff Theorem.

Proof. We have seen that \mathbb{R}_ℓ^2 is not normal. But since \mathbb{R}_ℓ is normal and therefore, completely regular, so is \mathbb{R}_ℓ^2 . Consequently, \mathbb{R}_ℓ^2 is homeomorphic to a subspace of $[0, 1]^J$ for some indexing set J . Due to the Tychonoff Theorem, $[0, 1]^J$ is compact Hausdorff, and thus, normal. This completes the proof. ■

3.5 The Urysohn Metrization Theorem

Theorem 3.37 (Urysohn Metrization Theorem). Every regular space X with a countable basis is metrizable.

Proof. Recall first that every regular space with a countable basis is normal. We shall show that X is metrizable by constructing an imbedding of X into \mathbb{R}^ω . We shall first show that there is a countable sequence of functions $\{f_n\}$ from X to $[0, 1]$ such that for all $x_0 \in X$ and a neighborhood U of x_0 , there is a function f_n such that $f_n(x_0) > 0$ and $f_n(x) = 0$ for all $x \in X \setminus U$.

Let $\mathcal{B} = \{B_n\}$ be a countable basis for X . Then, for all pairs (m, n) such that $\overline{B_m} \subseteq B_n$, define the function $g_{m,n} : X \rightarrow [0, 1]$, using Urysohn's Lemma, such that $g_{m,n}(\overline{B_m}) = \{1\}$ and $g_{m,n}(B_n) = \{0\}$. Obviously, the set $\{g_{m,n} \mid m, n \in \mathbb{N}\}$ is countable and it is not hard to see that this is our desired sequence of functions $\{f_n\}$.

From Theorem 3.34 and the sequence of functions constructed above, we see that X can be imbedded into \mathbb{R}^ω , and hence, is metrizable. ■

3.6 Tietze Extension Theorem

Lemma 3.38. Let X be a normal space and A be a closed subspace of X . Let $f : A \rightarrow [-r, r]$ be a continuous map. Then, there is a continuous function $g : X \rightarrow [-r, r]$ such that

$$|f(a) - g(a)| \leq 2r/3 \quad |g(x)| \leq r/3 \quad \text{for all } a \in A, x \in X$$

Proof. Define

$$I_1 = \left[-r, -\frac{r}{3}\right] \quad I_2 = \left[-\frac{r}{3}, \frac{r}{3}\right] \quad I_3 = \left[\frac{r}{3}, r\right]$$

and

$$B = f^{-1}(I_1) \quad C = f^{-1}(I_3)$$

Since I_1 and I_3 are closed in $[-r, r]$, B and C must be disjoint and closed in X . Now, due to Urysohn's Lemma, there is a function $g : X \rightarrow [-r/3, r/3]$ such that $g(B) = \{-r/3\}$ and $g(C) = \{r/3\}$, which has a natural extension $g : X \rightarrow [-r, r]$.

Obviously, $|g(x)| \leq r/3$ for all $x \in X$. Further, for all $a \in A$, if $a \in B$, then $g(a) = -r/3$, and $f(a) \in I_1$, similarly, if $a \in C$, then $g(a) = r/3$ and $f(a) \in I_3$. This immediately implies the desired conclusion. ■

Theorem 3.39 (Tietze Extension Theorem). *Let X be a normal space; let A be a closed subspace of X*

1. *Any continuous map of A into the closed interval $[-1, 1]$ of \mathbb{R} may be extended to a continuous map of all X into $[-1, 1]$*
2. *Any continuous map of A into \mathbb{R} may be extended to a continuous map of all of X into \mathbb{R}*

Proof. The main idea of the proof is to construct a uniformly convergent sequence of continuous functions to f on A . This would immediately imply the continuity of the limiting function over X , due to the Uniform Limit Theorem.

1. Using the preceding lemma, there is a function $g_1 : X \rightarrow [-1, 1]$ such that $|f(a) - g_1(a)| \leq 2/3$, while $|g_1(x)| \leq 1/3$ for all $a \in A$ and $x \in X$. Let us define $f_1 : A \rightarrow [-2/3, 2/3]$ as

$$f_1(x) = f(x) - g_1(x)$$

which is a continuous function. Then, we may reuse the previous lemma to define a function $g_2(x) : X \rightarrow [-1, 1]$ such that $|f_1(a) - g_2(a)| \leq (2/3)^2$, while $|g_2(x)| \leq (2/3)(1/3)$ and so on. As a result, we define the function $g_n : X \rightarrow [-1, 1]$ satisfying

$$|f_{n-1}(a) - g_n(a)| \leq \left(\frac{2}{3}\right)^n \quad |g_n(x)| \leq \frac{1}{3} \left(\frac{2}{3}\right)^{n-1}$$

Finally, define the functions $s_n : X \rightarrow \mathbb{R}$

$$s_n(x) = \sum_{i=1}^n g_i(x)$$

We note that

$$-1 < -\frac{1}{3} \sum_{i=1}^n \left(\frac{2}{3}\right)^{i-1} \leq s_n(x) \leq \frac{1}{3} \sum_{i=1}^n \left(\frac{2}{3}\right)^{i-1} < 1$$

Hence, we may take the restriction of s_n to $[-1, 1]$, which would also be continuous since it is the range restriction of a sum of finitely many continuous functions. Now, due to the Weierstrass M -test, the sequence of functions s_n are uniformly convergent. Further, since

$$|f(a) - s_n(a)| \leq \left(\frac{2}{3}\right)^n$$

we know that the convergent function $s : X \rightarrow [-1, 1]$ agrees with f on A . This completes the proof.

2. Recall that the spaces $(-1, 1)$ and \mathbb{R} are homeomorphic. Therefore, it suffices to prove the statement for functions of the form $f : A \rightarrow (-1, 1)$. Using the first part of this theorem, we know that there is a function $g : X \rightarrow [-1, 1]$. We shall use this function to obtain an extension h of f from $X \rightarrow (-1, 1)$. Let $D = g^{-1}(\{-1\}) \cup g^{-1}(\{1\})$. Since G is continuous, D is closed in X and must be disjoint from A . Then, using Urysohn's Lemma, there is a function $\phi : X \rightarrow [0, 1]$ such that $\phi(A) = \{1\}$ and $\phi(D) = \{0\}$. Then, the function $h(x) = \phi(x) \cdot g(x)$ is a continuous function from X to $(-1, 1)$ that agrees with f on A . This completes the proof. ■

Chapter 4

Nets and Filters

4.1 Nets

Definition 4.1 (Directed Set, Net, Cofinal). A *directed set* is a reflexive and transitive relation (I, \leq) such that for all $\alpha, \beta \in I$, there is $\gamma \in I$ such that $\alpha \leq \gamma$ and $\beta \leq \gamma$. A subset $J \subseteq I$ is said to be *cofinal* in I if for every $\alpha \in I$, there is $\beta \in J$ such that $\alpha \leq \beta$.

Proposition 4.2. If J is cofinal in (I, \leq) , then (J, \leq) is a directed set.

Proof. Obviously, (J, \leq) is a reflexive and transitive relation. Let $\alpha, \beta \in J$, then there is $\gamma \in I$ such that $\alpha, \beta \leq \gamma$. Now, there is $\delta \in J$ such that $\gamma \leq \delta$ and the conclusion follows. ■

Definition 4.3 (Net, Convergence). A *net* is a function from a directed set (I, \leq) to a topological space X . A net $(x_\alpha)_{\alpha \in I}$ is said to *converge* to $x \in X$ if for each neighborhood U of x , there is $\alpha \in I$ such that

$$\alpha \leq \beta \implies x_\beta \in U$$

Notice that the convergence in the case of nets reduces to the convergence of sequences when $(I, \leq) = (\mathbb{N}, \leq)$ with the standard total ordering.

Lemma 4.4. If X is Hausdorff, then a net in X converges to at most one point.

Proof. Suppose $(x_\alpha)_{\alpha \in I}$ converges to at least two distinct points x_1, x_2 . Then, there are disjoint neighborhoods U and V of x_1 and x_2 respectively. There is an index $\alpha \in I$ such that $x_t \in U$ whenever $\alpha \leq t$ and $\beta \in I$ such that $x_t \in V$ whenever $\beta \leq t$.

There is $\gamma \in I$ such that $\alpha, \beta \leq \gamma$, consequently, $x_\gamma \in U \cap V = \emptyset$, a contradiction. ■

Theorem 4.5. Let $A \subseteq X$. Then $x \in \overline{A}$ if and only if there is a net of point of A converging to x .

Proof. Suppose $x \in \overline{A}$. Let \mathcal{U} be the set of all neighborhoods of x . Define $U \leq V$ in \mathcal{U} if $U \supseteq V$. For each $U \in \mathcal{U}$, pick some $x_U \in U \cap A$. This is a net in X that converges to x . The converse is trivial. ■

Theorem 4.6. *A function $f : X \rightarrow Y$ is continuous if and only if it maps convergent nets to convergent nets.*

Proof. ■

Definition 4.7. Let $\mathbf{x} : (I, \leq) \rightarrow X$ be a net. If (K, \leq) is a directed set and $f : (K, \leq) \rightarrow (I, \leq)$ is an order preserving map such that $f(K)$ is cofinal in I , then $f \circ \mathbf{x} : K \rightarrow X$ is called a *subnet* of $(x_\alpha)_{\alpha \in I}$.

Proposition 4.8. *If $(x_\alpha)_{\alpha \in I}$ is a net converging to x , then every subnet converges to x .*

Proof. ■

Chapter 5

The Tychonoff Theorem

5.1 The Tychonoff Theorem

5.1.1 Filters and Ultrafilters

Definition 5.1 (Filter). A *filter* on a set X is a subset \mathcal{F} of $\mathcal{P}(X)$ satisfying:

- (a) $\emptyset \notin \mathcal{F}$
- (b) If $A \in \mathcal{F}$ and $A \subseteq B \subseteq X$, then $B \in \mathcal{F}$
- (c) If $A, B \in \mathcal{F}$, then $A \cap B \in \mathcal{F}$

It is not hard to see, using (a) and (b) that $X \in \mathcal{F}$ for every filter \mathcal{F} on X .

A subset \mathcal{A} of $\mathcal{P}(X)$ is said to have the *finite intersection property* if the intersection of a finite subset of \mathcal{A} is nonempty.

Lemma 5.2. Let $\mathcal{A} \subseteq \mathcal{P}(X)$ have the finite intersection property. Then there is a minimal filter containing \mathcal{A} .

Proof. Define \mathcal{U} to be the closure of \mathcal{A} under finite intersection. Next, define

$$\mathcal{F} = \{S \subseteq X \mid \exists A \in \mathcal{U}, A \subseteq S\}$$

We shall show that \mathcal{F} is a filter on X . Since \mathcal{U} does not contain \emptyset , neither does \mathcal{F} . Now, suppose $A \in \mathcal{F}$ and $A \subseteq B$, then by definition, there is $C \in \mathcal{U}$ such that $C \subseteq A$, consequently, $C \subseteq B$ and $B \in \mathcal{F}$. Finally, if $A, B \in \mathcal{F}$, then there are $C, D \in \mathcal{U}$ such that $C \subseteq A$ and $D \subseteq B$. Then, $C \cap D \in \mathcal{U}$ such that $C \cap D \subseteq A \cap B$, and thus, $A \cap B \in \mathcal{F}$. This shows that \mathcal{F} is a filter.

Let \mathcal{S} be the set of all filters on X containing \mathcal{A} . We know that \mathcal{S} is nonempty due to the above discussion. Now, define

$$\mathcal{F} = \bigcap_{\mathcal{F} \in \mathcal{S}} \mathcal{F}$$

from here it is not hard to see that \mathcal{F} is the unique minimal filter containing \mathcal{A} . ■

Definition 5.3 (Ultrafilter). An *ultrafilter* on X is a filter that is maximal with respect to the containment (partial) order.

Theorem 5.4. Let \mathcal{F} be a filter on X . Then \mathcal{F} is an ultrafilter if and only if for all $A \subseteq X$, either $A \in \mathcal{F}$ or $X \setminus A \in \mathcal{F}$ but not both.

Proof. Let \mathcal{F} be an ultrafilter on X and suppose $A \subseteq X$ such that $A \notin \mathcal{F}$ and $X \setminus A \notin \mathcal{F}$. Define $\mathcal{F}' = \mathcal{F} \cup \{A\}$. Then, $\mathcal{F}' \subseteq \mathcal{P}(X)$ and has the finite intersection property, consequently is contained in a filter $\overline{\mathcal{F}}$ due to Lemma 5.2 and hence, $\mathcal{F} \subsetneq \mathcal{F}' \subseteq \overline{\mathcal{F}}$, a contradiction to the maximality of \mathcal{F} .

Conversely, let \mathcal{F} be a filter on X satisfying the statement of the theorem. Suppose there is a filter \mathcal{F}' on X satisfying $\mathcal{F} \subsetneq \mathcal{F}'$. Let $A \in \mathcal{F}' \setminus \mathcal{F}$, then by the hypothesis, $X \setminus A \in \mathcal{F}$, and thus $X \setminus A \in \mathcal{F}'$, whence $\emptyset = A \cap (X \setminus A) \in \mathcal{F}'$, a contradiction. Hence, \mathcal{F} is an ultrafilter. ■

Lemma 5.5. Every filter is contained in an ultrafilter.

Proof. Let \mathcal{F} be a filter on a set X and \mathcal{S} be the set of all filters on X containing \mathcal{F} . Notice that \mathcal{S} forms a poset under containment. Let \mathcal{C} be a chain in the poset (\mathcal{S}, \subseteq) . Define the collection

$$\mathcal{F} = \bigcup_{F \in \mathcal{C}} F$$

We claim that \mathcal{F} is a filter on X . Indeed, $\emptyset \notin F$ for all $F \in \mathcal{C}$, whence $\emptyset \notin \mathcal{F}$. If $A, B \in \mathcal{F}$, then there is $F \in \mathcal{C}$ containing both A and B (since \mathcal{C} is a chain). Therefore, $A \cap B \in F \subseteq \mathcal{F}$. Finally, suppose $A \in \mathcal{F}$ and $A \subseteq B$, then there is $F \in \mathcal{C}$ such that $A \in F$, from which it would follow that $B \in F \subseteq \mathcal{F}$.

Finally, since \mathcal{F} is an element of \mathcal{S} , the chain \mathcal{C} is bounded above. Invoking Zorn's Lemma, we have a maximal element in \mathcal{S} with respect to inclusion, which is an ultrafilter containing \mathcal{F} . ■

Corollary 5.6. Let $\mathcal{A} \subseteq \mathcal{P}(X)$ have the finite intersection property. Then there is a ultrafilter on X containing \mathcal{A} .

Proposition 5.7. Let \mathcal{F} be an ultrafilter on X and $A \subseteq X$ such that A intersects every element in \mathcal{F} . Then, $A \in \mathcal{F}$.

Proof. Suppose $A \notin \mathcal{F}$. Then, due to Theorem 5.4, $X \setminus A \in \mathcal{F}$, a contradiction to the hypothesis that A intersects every element in \mathcal{F} . ■

Definition 5.8 (Filter Convergence). A filter \mathcal{F} on a topological space X is said to converge to $x \in X$ if for every neighborhood U of x , $U \in \mathcal{F}$.

Definition 5.9 (Pushforward). Let \mathcal{F} be a filter on X and $f : X \rightarrow Y$ be a map of sets. Then

$$f_*\mathcal{F} = \{A \subseteq Y \mid f^{-1}(A) \in \mathcal{F}\}$$

is a filter on Y .

Theorem 5.10. As defined above, $f_*\mathcal{F}$ is indeed a filter on Y . Further, if \mathcal{F} is an ultrafilter, then so is $f_*\mathcal{F}$

Proof. ■

5.1.2 First Proof

This version of the proof requires a more suitable characterization of compactness:

Theorem 5.11. *A topological space X is compact if and only if for every collection \mathcal{C} of closed sets in X having the finite intersection property,*

$$\bigcap_{C \in \mathcal{C}} C$$

is nonempty.

We are now ready to prove the Tychonoff Theorem.

Proof 1 of The Tychonoff Theorem. Let \mathcal{A} be a collection of subsets of $\prod_{\alpha \in J} X_{\alpha}$ having the finite intersection property. It suffices to show that

$$\bigcap_{A \in \mathcal{A}} \bar{A}$$

is nonempty.

Let \mathcal{F} be an ultrafilter containing \mathcal{A} . We shall show that

$$\bigcap_{F \in \mathcal{F}} \bar{F}$$

is nonempty, from which the result would follow.

Since \mathcal{F} has the finite intersection property, so does $\pi_{\alpha}(\mathcal{F}) = \{\pi_{\alpha}(F) \mid F \in \mathcal{F}\}$. Consequently, there is $x_{\alpha} \in X_{\alpha}$ such that $x_{\alpha} \in \pi_{\alpha}(F)$ for all $F \in \mathcal{F}$. Let $\mathbf{x} = (x_{\alpha})_{\alpha \in J}$. We claim that $\mathbf{x} \in \bar{F}$ for all $F \in \mathcal{F}$.

First, we shall show that every subbasis element containing \mathbf{x} intersects every element of \mathcal{F} . Consider the subbasis element $\pi_{\alpha}^{-1}(U_{\alpha})$ where U_{α} is a neighborhood of x_{α} in X_{α} . Since $x_{\alpha} \in \pi_{\alpha}(F)$ for all $F \in \mathcal{F}$, $U_{\alpha} \cap \pi_{\alpha}(F) \neq \emptyset$, consequently, $\pi_{\alpha}^{-1}(U_{\alpha}) \cap F \neq \emptyset$ for all $F \in \mathcal{F}$. From here, using Proposition 5.7, \mathcal{F} contains every subbasis element containing \mathbf{x} whence it contains every basis element containing \mathbf{x} .

Finally, let U be an open set in X containing \mathbf{x} . Then, U contains a basis element, say B that contains \mathbf{x} . Due to the preceeding paragraph, $B \in \mathcal{F}$, consequently, $B \cap F \neq \emptyset$ for all $F \in \mathcal{F}$ and therefore $U \cap F \neq \emptyset$ for all $F \in \mathcal{F}$ and $\mathbf{x} \in \bar{F}$ for all $F \in \mathcal{F}$. This completes the proof. ■

5.1.3 Second Proof

We first characterise compactness using the convergence of filters and ultrafilters.

Theorem 5.12 (Ultrafilter Convergence Theorem). *Let X be a topological space. X is compact if and only if every ultrafilter \mathcal{F} on X converges to at least one point.*

Proof. Suppose X is compact. Since \mathcal{F} has the finite intersection property, we must have $\bigcap_{F \in \mathcal{F}} \bar{F} \neq \emptyset$, and thus there is $x \in X$ such that $x \in \bar{F}$ for all $F \in \mathcal{F}$. Let U be a neighborhood of x in X . Then $U \cap F \neq \emptyset$ for all $F \in \mathcal{F}$, and due to Proposition 5.7, $U \in \mathcal{F}$. Thus \mathcal{F} converges to x .

Now, suppose X is not compact. Then, there is an open cover $\{U_{\alpha}\}_{\alpha \in J}$ that has no finite subcover. Define $A_{\alpha} = X \setminus U_{\alpha}$. Then $\{A_{\alpha}\}$ has the finite intersection property and hence is contained in an ultrafilter \mathcal{F} . Suppose \mathcal{F} does converge to a point $x \in X$. Choose $\beta \in J$ such that $x \in U_{\beta}$. By choice of x , we must have $U_{\beta} \in \mathcal{F}$, but this is a contradiction to $A_{\beta} \in \mathcal{F}$. ■

We are now ready to prove the Tychonoff Theorem.

Proof 2 of The Tychonoff Theorem. Let \mathcal{U} be an ultrafilter on X . Then $(\pi_\alpha)_* \mathcal{U}$ is an ultrafilter on X_α , consequently, converges to a point $x_\alpha \in X_\alpha$. Define the point $\mathbf{x} = (x_\alpha)_{\alpha \in J}$. We contend that \mathcal{U} converges to \mathbf{x} . Indeed, let U be an open set containing \mathbf{x} , then it contains a basis element $B = \prod_{\alpha \in J} U_\alpha$ containing \mathbf{x} . If we show that B is contained in \mathcal{U} , then it would immediately imply that U is contained in \mathcal{U} . But since each U_α is contained in $(\pi_\alpha)_* \mathcal{U}$, by definition, $\pi_\alpha^{-1}(U_\alpha)$ is contained in \mathcal{U} . Due to this and the fact that \mathcal{U} is closed under finite intersection, we have the desired conclusion. ■

5.2 Stone-Čech Compactification

Definition 5.13 (Compactification). A compactification of a space X is a compact Hausdorff space Y containing X as a subspace such that $\overline{X} = Y$. Two compactifications Y_1 and Y_2 of X are said to be equivalent if there is a homeomorphism $h : Y_1 \rightarrow Y_2$ such that $h(x) = x$ for all $x \in X$.

Lemma 5.14. Let X be a space and $h : X \rightarrow Z$ be an imbedding of X in the compact Hausdorff space Z . Then, there is a corresponding compactification Y of X such that there is an imbedding $H : Y \rightarrow Z$ that agrees with h on X . Further, the compactification Y is uniquely determined up to equivalence.

Y is called the **compactification induced** by the imbedding h .

Theorem 5.15. Let X be completely regular. Then there is a compactification Y of X having the property that every bounded continuous map $f : X \rightarrow \mathbb{R}$ extends uniquely to a continuous map of Y into \mathbb{R} .

Proof. Let $\{f_\alpha\}_{\alpha \in J}$ be the collection of all bounded continuous functions $f_\alpha : X \rightarrow \mathbb{R}$. Due to Theorem 3.34 and the fact that complete regularity implies the separation of points from open sets with a bounded function, there is an imbedding $F : X \rightarrow Z = \prod_{\alpha \in J} [\inf f_\alpha(X), \sup f_\alpha(X)]$. Due to the Tychonoff Theorem, the space Z is compact Hausdorff and hence, using Lemma 5.14, there is a compactification Y of X and a map $H : Y \rightarrow Z$ which agrees with F on X . Now, by taking the projection map π_α , we have a continuous map h that agrees with f_α on X .

The uniqueness of extension follows from a trivial property of a Hausdorff image space. ■

Theorem 5.16. Let X be completely regular and Y be a compactification of X satisfying the extension property of Theorem 5.15. Given any continuous map $f : X \rightarrow K$ to a compact Hausdorff space K , the map f extends uniquely to a continuous map $g : Y \rightarrow K$.

Proof. Since K is compact Hausdorff and therefore completely regular, it may be imbedded into $[0, 1]^J$ for some indexing set J . Let $f = (f_\alpha)$ be the imbedding. Then each map $f_\alpha : X \rightarrow [0, 1] \hookrightarrow \mathbb{R}$ may be extended uniquely to a map $f_\alpha : Y \rightarrow \mathbb{R}$. Further,

$$f_\alpha(Y) = f_\alpha(\overline{X}) \subseteq \overline{f_\alpha(X)} = [0, 1]$$

and thus, we have an extension $f : Y \rightarrow K$. This completes the proof. ■

The Universal Property

Fix some completely regular space X and consider the category \mathcal{C} of all maps $f : X \rightarrow K$ for some compact Hausdorff space K . A morphism between two objects f and g in \mathcal{C} is a continuous map h such that the

following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{g} & K_2 \\ f \downarrow & \nearrow h & \\ K_1 & & \end{array}$$

It is immediate from the definition of the category that the map $X \hookrightarrow Y$ is universal in this category and therefore is unique upto a unique isomorphism.

This object is known as the *Stone-Čech compactification* of X and is denoted by βX .

Chapter 6

Metrization and Paracompactness

Definition 6.1 (Locally Finite). Let X be a topological space. A collection \mathcal{A} of subsets of X is said to be *locally finite* in X if every point of X has a neighborhood that intersects only finitely many elements of \mathcal{A} .

Definition 6.2 (Countably Locally Finite). A collection \mathcal{B} of subsets of X is said to be *countably locally finite* if \mathcal{B} can be written as the countable union of collections \mathcal{B}_n , each of which is locally finite.

6.1 Paracompactness

Definition 6.3 (Paracompact). A topological space X is said to be *paracompact* if every open covering has a locally finite open refinement.

Proposition 6.4. *A paracompact Hausdorff space is regular.*

Proof. Let $A \subseteq X$ be a closed set and $x \notin A$. Then due to Hausdorff-ness, for each $a \in A$, there is an open neighborhood U_a of a whose closure does not contain x . Consider the indexed set $\{U_a\}_{a \in A} \cup \{X \setminus A\}$, which is an open cover of X whence has a locally finite open refinement, say \mathcal{U} . Let

$$\mathcal{V} = \{U \in \mathcal{U} \mid U \subseteq U_a \text{ for some } a \in A\}.$$

Then, \mathcal{V} is an open cover for A and for any $V \in \mathcal{V}$, there is some U_a such that $V \subseteq U_a$, whence $\overline{V} \subseteq \overline{U_a}$ and thus does not contain x .

Now, let

$$\tilde{V} = \bigcup_{V \in \mathcal{V}} V \quad \tilde{U} = X \setminus \underbrace{\left(\bigcup_{V \in \mathcal{V}} \overline{V} \right)}_{\text{since } \mathcal{V} \text{ is locally finite}} = X \setminus \left(\bigcup_{V \in \mathcal{V}} \overline{V} \right).$$

It is not hard to see that \tilde{U} and \tilde{V} form an open separation of x and A . ■

Proposition 6.5. *A paracompact Hausdorff space is normal.*

Proof. Proceed as above with a little tweaking. ■

Lemma 6.6. *Let X be regular. Then the following are equivalent:*

- (a) *Every open cover of X has a countably locally finite open refinement which is a cover.*
- (b) *Every open cover of X has a locally finite refinement which is a cover.*
- (c) *Every open cover of X has a locally finite closed refinement which is a cover.*
- (d) *Every open cover of X has a locally finite open refinement which is a cover.*

Proof. (a) \implies (b). Let \mathcal{A} be an open cover of X .

(c) \implies (d). This is the hardest part of the proof. Let \mathcal{A} be an open cover of X and let \mathcal{B} be a closed refinement of \mathcal{A} which is also a cover. Define

$$\mathcal{U} := \{U \subseteq_{\text{open}} X \mid U \text{ intersects finitely many elements in } \mathcal{B}\}.$$

Then \mathcal{U} is also an open cover of X whence has a locally finite closed refinement \mathcal{C} which is also a cover. Now, for each $B \in \mathcal{B}$, let

$$\mathcal{C}(B) := \{C \mid C \in \mathcal{C} \text{ and } C \cap B = \emptyset\}.$$

and let

$$E(B) := X \setminus \bigcup_{C \in \mathcal{C}(B)} C.$$

Then $E(B)$ is an open¹ set containing B . For each $B \in \mathcal{B}$, pick $F(B) \in \mathcal{A}$ containing B and finally define

$$\mathcal{D} := \{E(B) \cap F(B) \mid B \in \mathcal{B}\}.$$

That this is an open refinement of \mathcal{A} covering X is trivial. It remains to show that \mathcal{D} is locally finite. Let $x \in X$. Then, there is a neighborhood W of x that intersects finitely elements in \mathcal{C} , say C_1, \dots, C_k and since \mathcal{C} forms a cover, $W \subseteq C_1 \cup \dots \cup C_k$.

Note that if $C \in \mathcal{C}$ intersecting $E(B) \cap F(B)$ for some $B \in \mathcal{B}$, then C intersects $E(B)$ whence C intersects B by definition. Owing to the local finiteness of \mathcal{B} , C intersects B and hence C can intersect only finitely many elements of \mathcal{D} .

Whence W can intersect only finitely many elements of \mathcal{D} implying the desired conclusion.

(d) \implies (a). Trivial. This completes the proof. ■

Proposition 6.7 (A. H. Stone). *Every metrizable space is paracompact.*

The machinery and proof is due to M. E. Rudin.

Proof. ■

Proposition 6.8. *A regular Lindelöf space is paracompact.*

Proof. Let \mathcal{A} be an open cover of a regular Lindelöf space X . Then, there is a countable subcollection that covers X . Since we have a countably locally finite refinement of \mathcal{A} which is a cover, due to Lemma 6.6, the space is paracompact. ■

¹Since $C(B)$ is locally finite.

6.1.1 Partitions of Unity

Definition 6.9 (Partition of Unity). Let $\{U_\alpha\}_{\alpha \in J}$ be an indexed open covering of X . An indexed family of continuous functions $\phi_\alpha : X \rightarrow [0, 1]$ is said to be a *partition of unity on X dominated by $\{U_\alpha\}_{\alpha \in J}$* if

- (a) $\text{Supp } \phi_\alpha \subseteq U_\alpha$ for each $\alpha \in J$.
- (b) $\{\text{Supp } \phi_\alpha\}_{\alpha \in J}$ is locally finite.
- (c) $\sum_{\alpha \in J} \phi_\alpha(x) = 1$ for each $x \in X$.

Lemma 6.10 (Shrinking Lemma). Let X be a paracompact Hausdorff space and $\{U_\alpha\}_{\alpha \in J}$ be an indexed open cover of X . Then there is a locally finite indexed open cover $\{V_\alpha\}_{\alpha \in J}$ of X such that $\overline{V_\alpha} \subseteq U_\alpha$ for each $\alpha \in J$.

Proof. Let \mathcal{A} denote the collection of all open sets U in X such that \overline{U} is contained in some U_α . We first contend that \mathcal{A} forms an open cover of X . Indeed, every $x \in X$ is contained in some U_α and since X is regular, there is a neighborhood V of x such that $x \in V \subseteq \overline{V} \subseteq U_\alpha$ whence x is covered by \mathcal{A} .

Since X is paracompact, the open cover \mathcal{A} has a locally finite open refinement \mathcal{B} . For each $\alpha \in J$, define

$$V_\alpha = \bigcup_{\substack{B \in \mathcal{B} \\ B \subseteq U_\alpha}} B \subseteq U_\alpha$$

Since the set $\{B \in \mathcal{B} \mid B \subseteq U_\alpha\} \subseteq \mathcal{B}$ is locally finite, we have

$$\overline{V_\alpha} = \bigcup_{\substack{B \in \mathcal{B} \\ B \subseteq U_\alpha}} \overline{B} \subseteq U_\alpha.$$

Further, using the fact that X is regular, it is not hard to see that if $x \in U_\alpha$, then $x \in V_\alpha$ and thus the collection $\{V_\alpha\}_{\alpha \in J}$ forms an open cover for X and is an open refinement of $\{U_\alpha\}_{\alpha \in J}$.

Lastly, we must show that $\{V_\alpha\}$ is locally finite. Let $x \in X$. Since $\{U_\alpha\}$ was locally finite, there is a neighborhood U of x which intersects finitely many of the U_α 's whence it intersects finitely many of the V_α 's. This completes the proof. \blacksquare

Theorem 6.11. Let X be a paracompact Hausdorff space and $\{U_\alpha\}_{\alpha \in J}$ an indexed open cover of X . Then there is a partition of unity on X dominated by $\{U_\alpha\}_{\alpha \in J}$.

Proof. First, by repeated application of Lemma 6.10, there are successive locally finite precise open refinements $\{W_\alpha\}_{\alpha \in J}$ and $\{V_\alpha\}_{\alpha \in J}$ such that for each $\alpha \in J$,

$$\overline{W_\alpha} \subseteq V_\alpha \quad \text{and} \quad \overline{V_\alpha} \subseteq U_\alpha.$$

Using Lemma 3.21, there is a continuous function $\phi_\alpha : X \rightarrow [0, 1]$, such that $\phi_\alpha(\overline{W_\alpha}) = 1$ and $\phi_\alpha(X \setminus V_\alpha) = 0$. It is not hard to see that

$$\text{Supp } \phi_\alpha \subseteq \overline{V_\alpha} \subseteq U_\alpha.$$

Consider the function $\Phi : X \rightarrow \mathbb{R}$ given by

$$\Phi(x) = \sum_{\alpha \in J} \phi_\alpha(x).$$

Let $x \in X$. Due to local finiteness, there is an open neighborhood U of x such that U intersects only finitely many of the V'_α 's whence $U \subseteq X \setminus V_\alpha$ for all but finitely many $\alpha \in J$, whence the sum makes sense because it is essentially finite.

Moreover, since $\{\overline{W_\alpha}\}$ is an open cover of X , every $x \in X$ belongs to at least one $\overline{W_\alpha}$ and thus $\Phi(x)$ is positive for all $x \in X$. It is not hard to see now that the collection

$$\left\{ \frac{\phi_\alpha}{\Phi} \right\}_{\alpha \in J}$$

is a partition of unity dominated by $\{U_\alpha\}$. ■

Chapter 7

Complete Metric Spaces and Function Spaces

7.1 Complete Metric Spaces

Definition 7.1 (Completeness). A metric space is said to be complete if every Cauchy sequence converges.

Lemma 7.2. A metric space X is complete if and only if every Cauchy sequence has a convergent subsequence.

Proof. One direction is trivial. Suppose now that every Cauchy sequence has a convergent subsequence. Let $\{x_n\}$ be a Cauchy sequence and $\{x_{n_k}\}$ be the convergent subsequence that converges to $x \in X$. Let $\varepsilon > 0$. Then there is $K \in \mathbb{N}$ such that for all $k \geq K$, $d(x_{n_k}, x) < \varepsilon/2$. Further, since the sequence is Cauchy, there is $N \in \mathbb{N}$ such that for all $m, n \geq N$, $d(x_m, x_n) < \varepsilon/2$. Let $l \geq K$ be such that $n_l \geq N$. As a result, for all $m \geq n_l$,

$$d(x_m, x) \leq d(x_m, x_{n_l}) + d(x_{n_l}, x) < \varepsilon$$

This completes the proof. ■

Corollary 7.3. A compact metric space is complete. Further, \mathbb{R}^n is complete for each $n \in \mathbb{N}$.

Proof. The first assertion follows from the fact that every compact metric space is sequentially compact. Since every Cauchy sequence is bounded, it is contained in some closed n -cell, which is compact. The conclusion follows. ■

Lemma 7.4. Let $X = \prod_{\alpha \in J} X_\alpha$ in the product topology. Let \mathbf{x}_n be a sequence of points of X . Then $\mathbf{x}_n \rightarrow \mathbf{x}$ if and only if $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$.

Proof. The forward direction follows trivially from the fact that $\pi_\alpha : X \rightarrow X_\alpha$ is a continuous function. Conversely, suppose $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$ for each $\alpha \in J$. Let U be an open set containing \mathbf{x} , then U contains a basis element of the form $\prod_{\alpha \in J} U_\alpha$ where $U_\alpha = X_\alpha$ for all but finitely many $\alpha \in J$, say $\alpha_1, \dots, \alpha_k$. For each α_i , there is $N_i \in \mathbb{N}$ such that for all $n \geq N_i$, $\pi_{\alpha_i}(\mathbf{x}_n) \in U_{\alpha_i}$. Finally, letting $N = \max\{N_1, \dots, N_k\}$, we have that $\mathbf{x}_n \in \prod_{\alpha \in J} U_\alpha$ for all $n \geq N$. ■

Note that the above lemma is obviously not true for the box topology. Indeed, consider the sequence $\left\{ \frac{1}{n} \times \frac{1}{n} \times \cdots \right\}_{n \in \mathbb{N}}$ in \mathbb{R}^ω with the box topology. This sequence obviously does not converge to $\mathbf{0}$, but each component does converge to 0.

Theorem 7.5. *If the space Y is complete in the metric d , then the space Y^J is complete in the uniform metric $\bar{\rho}$ induced by d where J is any indexing set.*

Proof. Let $\varepsilon > 0$ be given. Set $\delta = \varepsilon/3$. There is $N \in \mathbb{N}$ such that for all $m, n \geq N$, $\bar{\rho}(f_m, f_n) < \delta$, consequently, $\bar{d}(f_m(\alpha), f_n(\alpha)) < \delta$ for each $\alpha \in J$. Using the completeness of Y , the sequence $\{f_n(\alpha)\}$ converges to say $f(\alpha)$ where $f : J \rightarrow Y$. Now, for each $\alpha \in J$, there is M_α such that for all $m \geq M_\alpha$, $\bar{d}(f_m(\alpha), f(\alpha)) < \delta$. As a consequence, $\bar{d}(f_m(\alpha), f_n(\alpha)) < 2\delta$ for each $n \geq N$ and thus $\bar{\rho}(f, f_n) \leq 2\delta < \varepsilon$. This completes the proof. ■

Proposition 7.6. *Let (X, d) be a complete metric space and $A \subseteq X$ be closed. Then A is complete.*

Proof. Let $\{a_n\}$ be a Cauchy sequence in A . Then, it is Cauchy in X , consequently, converges to a point $x \in X$. Since A is closed, $x \in \bar{A} = A$, and the conclusion follows. ■

Definition 7.7. Let X be a topological space and Y a metric space. Then $\mathcal{C}(X, Y)$ denotes the subspace of continuous functions from X to Y of Y^X and $\mathcal{B}(X, Y)$ denotes the subspace of bounded functions from X to Y of Y^X .

Theorem 7.8. *If the space Y is complete in the metric d and X a topological space, then Y^X , $\mathcal{C}(X, Y)$ and $\mathcal{B}(X, Y)$ are complete in the uniform metric corresponding to d .*

Proof. We shall show that $\mathcal{C}(X, Y)$ and $\mathcal{B}(X, Y)$ are closed in Y^X . The closedness of $\mathcal{C}(X, Y)$ follows from the uniform limit theorem. To show that $\mathcal{B}(X, Y)$ is closed, we shall show that its complement is open. Let $f : X \rightarrow Y$ be an unbounded function. We shall show that $B_{\bar{\rho}}(f, 1/2)$ is disjoint from $\mathcal{B}(X, Y)$. Let $g \in B_{\bar{\rho}}(f, 1/2)$. Fix some basepoint $y_0 \in Y$ and $M > 0$. There is $x \in X$ such that $d(y_0, f(x)) > M + 1/2$. As a result, $d(y_0, g(x)) > M$, this completes the proof. ■

7.2 Completion of a Metric Space

Theorem 7.9. *Let (X, d) be a metric space. Then there is a complete metric space (Y, D) and an isometric map $\Phi : (X, d) \rightarrow (Y, D)$ such that $\overline{\Phi(X)} = Y$. In this case, Y is called the **completion** of X .*

We present two proofs of this theorem. The first is inspired by the construction of the real numbers from the rationals, while the second imbeds X in a complete function space.

Proof 1. Let \tilde{X} denote the set of all Cauchy sequences in X . Since every constant sequence is Cauchy, the set \tilde{X} is nonempty. Consider the relation \sim on \tilde{X} , given by

$$\mathbf{x} \sim \mathbf{y} \iff \lim_{n \rightarrow \infty} d(x_n, y_n) = 0$$

We claim that \sim is an equivalence relation. The reflexivity and symmetry of \sim is obvious. It remains to show the transitivity of \sim . Indeed, if $\mathbf{x} \sim \mathbf{y}$ and $\mathbf{y} \sim \mathbf{z}$, then

$$0 \leq \lim_{n \rightarrow \infty} d(x_n, z_n) \leq \lim_{n \rightarrow \infty} d(x_n, y_n) + d(y_n, z_n) = 0$$

and the conclusion follows.

Let $Y = \tilde{X} / \sim$. Define the function $D : Y \times Y \rightarrow \mathbb{R}$ by

$$D([\mathbf{x}], [\mathbf{y}]) = \lim_{n \rightarrow \infty} d(x_n, y_n)$$

We must first show that D is well defined. Indeed, let $\mathbf{x}, \mathbf{x}' \in [\mathbf{x}]$ and $\mathbf{y}, \mathbf{y}' \in [\mathbf{y}]$. Then,

$$\lim_{n \rightarrow \infty} d(x'_n, y'_n) \leq \lim_{n \rightarrow \infty} d(x'_n, x_n) + d(x_n, y_n) + d(y_n, y'_n) = \lim_{n \rightarrow \infty} d(x_n, y_n)$$

and by symmetry, the inequality also holds in the other direction and thus D is well defined. We now show that D is a metric. To do so, notice that it suffices to verify that it satisfies the triangle inequality. Let $[\mathbf{x}], [\mathbf{y}], [\mathbf{z}] \in Y$. Then,

$$\lim_{n \rightarrow \infty} d(x_n, z_n) \leq \lim_{n \rightarrow \infty} d(x_n, y_n) + d(y_n, z_n) = D([\mathbf{x}], [\mathbf{y}]) + D([\mathbf{y}], [\mathbf{z}])$$

Consider the map $\Phi : (X, d) \rightarrow (Y, D)$ by $\Phi(x) = (x, x, \dots)$. It is obvious that this is an isometric imbedding. Let $[\mathbf{y}] \in Y$ where $\mathbf{y} = (y_n)$. Define $\mathbf{x}_n = (y_n, y_n, \dots)$. Since \mathbf{y} is a Cauchy sequence, it is not hard to see that $[\mathbf{x}_n] \rightarrow [\mathbf{y}]$. Hence, $\Phi(X)$ is dense in Y .

Next, note that any Cauchy sequence in $\Phi(X)$ converges in Y , for if $\{(x_n, x_n, \dots)\}$ is a Cauchy sequence in $\Phi(X)$, then it converges to $[(x_1, x_2, \dots)]$.

Finally, we shall show that Y is complete. Let $\{[\mathbf{y}_n]\}$ be a Cauchy sequence in Y . Since $\Phi(X)$ is dense in Y , there is a sequence $\{x_n\}$ in X such that $D(\Phi(x_n), [\mathbf{y}_n]) < 1/n$. It is not hard to see that $\{\Phi(x_n)\}$ is a Cauchy sequence and converges to some point $\mathbf{x} \in Y$. We contend that $[\mathbf{y}_n] \rightarrow \mathbf{x}$. Let $\varepsilon > 0$ be given and $N \in \mathbb{N}$ such that $1/N < \varepsilon/2$. Further, let $M \in \mathbb{N}$ such that for all $n \geq M$, $D(\Phi(x_n), \mathbf{x}) < \varepsilon/2$. As a result, for all $n \geq \max\{M, N\}$,

$$D([\mathbf{y}_n], \mathbf{x}) \leq D([\mathbf{y}_n], \Phi(x_n)) + D(\Phi(x_n), \mathbf{x}) < \varepsilon$$

This completes the proof. ■

Then next proof is shorter but less insightful.

Proof 2. We have already shown that $\mathcal{B}(X, \mathbb{R})$ is complete under the sup-metric. We shall now imbed (X, d) in $\mathcal{B}(X, \mathbb{R})$. Fix some point $x_0 \in X$. For each $a \in X$, define the function $\phi_a : X \rightarrow \mathbb{R}$ by

$$\phi_a(x) = d(x, a) - d(x, x_0)$$

It is not hard to see, using the triangle inequality that $|\phi_a(x)| \leq d(a, x_0)$. Thus, $\phi_a \in \mathcal{B}(X, \mathbb{R})$. We claim that the map $\Phi : X \rightarrow \mathcal{B}(X, \mathbb{R})$ given by $\Phi(a) = \phi_a$ is an imbedding under the sup-metric on $\mathcal{B}(X, \mathbb{R})$. Note that we have

$$|\phi_a(x) - \phi_b(x)| = |d(x, a) - d(x, x_0) - d(x, b) + d(x, x_0)| = |d(x, a) - d(x, b)| \leq d(a, b)$$

as a result, $\rho(\phi_a, \phi_b) \leq d(a, b)$. On the other hand, since $|\phi_a(a) - \phi_b(a)| = d(a, b)$, we must have $\rho(\phi_a, \phi_b) = d(a, b)$. Thus, Φ is an isometric imbedding. The conclusion follows by taking $Y = \overline{\Phi(X)}$. ■

7.3 Compactness in Metric Spaces

Definition 7.10 (Total Boundedness). A metric space (X, d) is said to be totally bounded if for each $\varepsilon > 0$, there is a finite cover of X by ε -balls.

Theorem 7.11. *A metric space (X, d) is compact if and only if it is complete and totally bounded.*

Proof. The forward direction is trivial. Conversely, suppose (X, d) is complete and totally bounded. We shall show that it is sequentially compact. Let $\{x_n\}$ be sequence in X . Using total boundedness, there is a covering of X with 1-balls. Choose the ball containing x_n for infinitely many indices n . Call the set of all such x_n 's as J_1 . Inductively, given J_{n-1} , consider a finite covering of X with $1/n$ -balls and choose the ball containing $x_k \in J_{n-1}$ for infinitely many indices k . Then $J_1 \supseteq J_2 \supseteq \dots$ and furthermore, $\text{diam } J_k \leq 2/k$. From each J_k , choose x_{n_k} such that $n_k > n_{k-1}$ which can obviously be done since each J_k contains x_n for infinitely many indices n .

Finally, it is not hard to see that $\{x_{n_k}\}$ forms a Cauchy sequence and is thus convergent. Hence (X, d) is sequentially compact, as a result, compact. ■

Definition 7.12 (Equicontinuous). Let X be a topological space and (Y, d) a metric space. A subset \mathcal{F} of $\mathcal{C}(X, Y)$ is said to be equicontinuous at $x_0 \in X$ if for each $\varepsilon > 0$, there is a neighborhood U of x_0 such that for all $x \in U$ and $f \in \mathcal{F}$, $d(f(x), f(x_0)) < \varepsilon$. Further, \mathcal{F} is said to be equicontinuous if it is equicontinuous at each point $x \in X$.

Theorem 7.13. *Let X be a topological space and (Y, d) a metric space. Then if $\mathcal{F} \subseteq \mathcal{C}(X, Y)$ is totally bounded under the uniform metric corresponding to d , then it is equicontinuous.*

Proof. Let $0 < \varepsilon < 1$ be given and $x_0 \in X$. Let $\delta = \varepsilon/3$. Then there is a collection $\{B_{\bar{\rho}}(f_i, \delta)\}_{i=1}^n$ of δ -balls that cover \mathcal{F} . Since this collection is finite, there is a neighborhood U of x_0 such that for all $x \in U$, $d(f_i(x), f_i(x_0)) < \delta$ for $1 \leq i \leq n$.

Let $f \in \mathcal{F}$. Then there is an index j such that $f \in B_{\bar{\rho}}(f_j, \delta)$ and hence $\bar{\rho}(f, f_j) < \delta$. Using the triangle inequality, we have, for all $x \in U$,

$$d(f(x), f(x_0)) < d(f(x), f_j(x)) + d(f_j(x), f_j(x_0)) + d(f_j(x_0), f(x_0)) < 3\delta = \varepsilon$$

This completes the proof. ■

Lemma 7.14. *Let X be a compact topological space and Y a compact metric space. If the subset \mathcal{F} of $\mathcal{C}(X, Y)$ is equicontinuous under d , then \mathcal{F} is totally bounded under the uniform and sup metrics corresponding to d .*

Proof. Since X is compact, all functions in $\mathcal{C}(X, Y)$ are bounded and as a result, the sup metric is well defined. We shall show total boundedness in the sup metric which would immediately imply total boundedness in the uniform metric.

Let $\varepsilon > 0$ and $\delta = \varepsilon/3$. Using equicontinuity, for each $x \in X$, there is a neighborhood U_x of x such that for all $t \in U_x$, $d(f(t), f(x)) < \delta$. Since X is compact, there is a finite cover, $\{U_{a_1}, \dots, U_{a_k}\}$. Using compactness and therefore, total boundedness of Y , there is a covering of Y by $\delta/2$ -balls, $\{V_1, \dots, V_m\}$.

Let J be the collection of all functions $\alpha : \{1, \dots, k\} \rightarrow \{1, \dots, m\}$ such that for there is a function $f_\alpha \in \mathcal{F}$ such that $f_\alpha(a_i) \in V_{\alpha(i)}$. We shall show that $\{B_{\bar{\rho}}(f_\alpha, \varepsilon)\}_{\alpha \in J}$ forms a cover for \mathcal{F} .

Let $f \in \mathcal{F}$. For each $1 \leq i \leq k$, choose an integer $\alpha(i) \in \{1, \dots, m\}$ such that $f(a_i) \in V_{\alpha(i)}$. Obviously, $\alpha \in J$ and $f_\alpha(a_i) \in V_{\alpha(i)}$. As a result, for all $x \in X$, choose a_i such that $x \in U_{a_i}$, then

$$d(f(x), f_\alpha(x)) \leq d(f(x), f(a_i)) + d(f(a_i), f_\alpha(a_i)) + d(f_\alpha(a_i), f_\alpha(x)) < 3\delta = \varepsilon$$

This completes the proof. ■

Definition 7.15. If (Y, d) is a metric space, a subset \mathcal{F} of $\mathcal{C}(X, Y)$ is said to be pointwise bounded under d if for each $x \in X$, the subset $\mathcal{F}_x = \{f(x) \mid f \in \mathcal{F}\}$ of Y is bounded under d .

Theorem 7.16 (Classical Ascoli's Theorem). Let X be a compact space and (\mathbb{R}^n, d) have the standard metric topology and give $\mathcal{C}(X, \mathbb{R}^n)$ the corresponding uniform topology. A subspace \mathcal{F} of $\mathcal{C}(X, \mathbb{R}^n)$ has compact closure if and only if \mathcal{F} is equicontinuous and bounded.

Proof. Since X is compact, the uniform metric and sup metric induce the same topologies. As a result, it suffices to consider the sup metric. Let \mathcal{G} denote the closure of \mathcal{F} in $\mathcal{C}(X, \mathbb{R}^n)$.

(\Rightarrow) Suppose \mathcal{G} is compact and therefore, is totally bounded under ρ . Using a preceding lemma, we may conclude that it is equicontinuous. Pointwise boundedness follows from the fact that it is compact and therefore bounded, as a result, $d(f(x), g(x)) < M$ for some $M > 0$ for all $f, g \in \mathcal{G}$ and $x \in X$. As a result, $\text{diam } \mathcal{G}_x \leq M$.

(\Leftarrow) Now suppose \mathcal{F} is equicontinuous and pointwise bounded. We shall first show that \mathcal{G} is equicontinuous and pointwise bounded. Since \mathcal{G} is the closure of \mathcal{F} , $\text{diam } \mathcal{G} = \text{diam } \mathcal{F}$ and pointwise boundedness follows. We now show equicontinuity. Let $\varepsilon > 0$ and choose some $x_0 \in X$. There is a neighborhood U of x_0 such that for all $x \in U$, $d(f(x), f(x_0)) < \varepsilon/3$. Let $g \in \mathcal{G}$. Then, there is some $f \in \mathcal{F}$ such that $\rho(f, g) < \varepsilon/3$. Then,

$$d(g(x), g(x_0)) \leq d(g(x), f(x)) + d(f(x), f(x_0)) + d(f(x_0), g(x_0)) < \varepsilon$$

for all $x \in U$. This shows equicontinuity.

Next, we shall show that $\bigcup_{g \in \mathcal{G}} g(X)$ is bounded in Y . Using equicontinuity, for each $a \in X$, there is a neighborhood U_a of a such that for all $x \in U_a$, and $g \in \mathcal{G}$, $d(g(x), g(a)) < 1$. Since X is compact, there is a finite cover $\{U_{a_1}, \dots, U_{a_k}\}$ of X . Now, for any $g \in \mathcal{G}$ and $x_1, x_2 \in X$, there are open sets, say U_{a_1} and U_{a_2} containing x_1 and x_2 respectively. Then,

$$\begin{aligned} d(g(x_1), g(x_2)) &\leq d(g(x_1), g(a_1)) + d(g(a_1), g(a_2)) + d(g(a_2), g(x_2)) \\ &< 2 + d(g(a_1), g(a_2)) \\ &\leq 2 + \max_{1 \leq i, j \leq k} d(g(a_i), g(a_j)) \end{aligned}$$

The last quantity is bounded since \mathcal{G}_{a_i} is bounded for each $1 \leq i \leq k$.

Finally, we shall show that \mathcal{G} is compact. Note that since it is a closed subset of $\mathcal{C}(X, \mathbb{R}^n)$, it is complete, therefore, it suffices to show that it is totally bounded. In the previous paragraph, we have shown that $\bigcup_{g \in \mathcal{G}} g(X)$ is bounded in \mathbb{R}^n and therefore, is a subset of a compact subspace Y of \mathbb{R}^n , consequently, we may view \mathcal{G} as a subspace of $\mathcal{C}(X, Y)$ and total boundedness follows from the previous lemma. ■

Theorem 7.17 (Classical Arzelà's Theorem). Let X be compact and let $f_n \in \mathcal{C}(X, \mathbb{R}^k)$ for all $k \in \mathbb{N}$. If the collection $\{f_n\}_{n=1}^\infty$ is pointwise bounded and equicontinuous, then it has a uniformly convergent subsequence.

Proof. Let $\mathcal{F} = \{f_n\}_{n=1}^\infty$ then due to Ascoli's Theorem, \mathcal{F} has compact closure \mathcal{G} in $\mathcal{C}(X, \mathbb{R}^k)$. Since \mathcal{G} is a compact metric space, it is sequentially compact, as a result, there is a subsequence $\{f_{n_j}\}$ which is convergent in the uniform metric, and thus uniformly convergent. ■

The following is the version of Arzelà-Ascoli that Rudin states:

Corollary 7.18. Let X be compact and $\{f_n\}$ be a sequence of functions in $\mathcal{C}(X, \mathbb{R}^k)$ for some positive integer k . If $\{f_n\}$ is pointwise bounded and equicontinuous on X , then

- (a) $\{f_n\}$ is uniformly bounded on X
- (b) $\{f_n\}$ contains a uniformly convergent subsequence

7.4 The Stone-Weierstrass Theorem

Definition 7.19 (Algebra). Let X be a topological space. A family $\mathcal{A} \subseteq \mathcal{C}(X, \mathbb{R})$ is said to be a real algebra if for all $f, g \in \mathcal{A}$ and $c \in \mathbb{R}$,

$$f + g \in \mathcal{A}, \quad fg \in \mathcal{A}, \quad cf \in \mathcal{A}$$

Similarly, a complex algebra satisfies all the above requirements along with $c \in \mathbb{C}$ and $\mathcal{A} \subseteq \mathcal{C}(X, \mathbb{C})$.

Definition 7.20 (Uniformly Closed). An algebra $\mathcal{A} \subseteq \mathcal{C}(X, \mathbb{K})$, where $\mathbb{K} = \mathbb{R}, \mathbb{C}$ is said to be uniformly closed if it is a closed set in the uniform topology given to $\mathcal{C}(X, \mathbb{K})$. Analogously define the uniform closure.

Definition 7.21. Suppose \mathcal{A} is an algebra of functions on a set X , then \mathcal{A} is said to separate points in X if for every pair of distinct points $x_1, x_2 \in X$, there is a function $f \in \mathcal{A}$ such that $f(x_1) \neq f(x_2)$.

If for each $x \in X$, there is a function $f \in \mathcal{A}$ such that $f(x) \neq 0$, then we say that \mathcal{A} vanishes at no point of E .

Lemma 7.22. Suppose \mathcal{A} is an algebra over a topological space X which separates points and vanishes at no point on X . Suppose $x_1, x_2 \in X$ are distinct and $c_1, c_2 \in \mathbb{C}$. Then there is $f \in \mathcal{A}$ such that $f(x_1) = c_1$ and $f(x_2) = c_2$.

Proof. According to the hypothesis on \mathcal{A} , there are continuous functions $g, h, k \in \mathcal{A}$ such that $g(x_1) \neq g(x_2)$ and $h(x_1) \neq 0$ and $k(x_2) \neq 0$. Define the functions

$$u = (g - g(x_1))k \quad v = (g - g(x_2))h$$

Since \mathcal{A} forms an algebra, $u, v \in \mathcal{A}$. Further, $u(x_1) = 0, u(x_2) \neq 0, v(x_1) \neq 0$ and $v(x_2) = 0$. Finally, define the function

$$f = c_1 \frac{v}{v(x_1)} + c_2 \frac{u}{u(x_2)}$$

Again, since \mathcal{A} forms an algebra, $f \in \mathcal{A}$ and the proof is complete. ■

Theorem 7.23 (Stone-Weierstrass Theorem). Let X be a compact Hausdorff space and $\mathcal{A} \subseteq \mathcal{C}(X, \mathbb{R})$. If \mathcal{A} separates points on X and vanishes at no point of X , then \mathcal{A} is dense in $\mathcal{C}(X, \mathbb{R})$ in the sup metric (which is the same as the uniform topology).

Proof. The proof proceeds in multiple steps. Let \mathcal{B} denote the closure of \mathcal{A} in $\mathcal{C}(X, \mathbb{R})$. Then, we have

Claim 1. If $f \in \mathcal{B}$, then $|f| \in \mathcal{B}$.

Proof. First, let $a = \sup_{x \in X} |f|$. Thus, $-a \leq f \leq a$ on X . Using the Weierstrass Approximation Theorem, there is a sequence of polynomials $\{p_n(t)\}$ that converge uniformly to $|t|$ on the interval $[-a, a]$. Note that the function $p_n \circ f$ is in the algebra \mathcal{A} , since it is a polynomial in f , therefore, the sequence $\{p_n(f)\}$ of continuous functions in \mathcal{A} converge in the sup metric to $|f|$, as a result, $|f| \in \mathcal{B}$, since the latter is a closed set.

Claim 2. If $f, g \in \mathcal{B}$ then $\max\{f, g\}, \min\{f, g\} \in \mathcal{B}$.

Proof. Follows from the following identities:

$$\max\{f, g\} = \frac{f + g + |f - g|}{2} \quad \min\{f, g\} = \frac{f + g - |f - g|}{2}$$

Claim 3. Let $f \in \mathcal{C}(X, \mathbb{R})$, $x \in X$ and $\varepsilon > 0$. Then there is a function $g_x \in \mathcal{B}$ such that $g_x(x) = f(x)$ and $g_x(t) > f(t) - \varepsilon$ for each $t \in X$.

Proof. Note that $\mathcal{A} \subseteq \mathcal{B}$ and due to Lemma 7.22, for every $y \in X$, there is a continuous function $h_y \in \mathcal{A}$ such that $h_y(x) = f(x)$ and $h_y(y) = f(y)$. Let U_y be the open set $(f - h_y)^{-1}((-\infty, \varepsilon))$. Notice that $\{U_y\}$ is an open cover for X , and thus has a finite subcover, say $\{U_{y_1}, \dots, U_{y_n}\}$. Finally, define the function

$$g_x = \max\{h_{y_1}, \dots, h_{y_n}\}$$

Note that in fact, $g_x \in \mathcal{A}$. The conclusion follows.

Claim 4. Let $f \in \mathcal{C}(X, \mathbb{R})$ and $\varepsilon > 0$, there exists a function $h \in \mathcal{A}$ such that $|h(x) - f(x)| < \varepsilon$

Proof. In the previous claim, we have constructed the functions g_x . Let $U_x = (g_x - f)^{-1}((-\infty, \varepsilon))$. Then, $\{U_x\}$ forms an open cover for X , and therefore has a finite subcover, say $\{U_{x_1}, \dots, U_{x_n}\}$. Define the function

$$h = \min\{g_{x_1}, \dots, g_{x_n}\}$$

It is not hard to see that $|h - f| < \varepsilon$ on X .

From the four claims, we see that for each $\varepsilon > 0$ and each $f \in \mathcal{C}(X, \mathbb{R})$, there is $h \in \mathcal{A}$ such that $|h - f| < \varepsilon$ and the conclusion follows. ■

Definition 7.24. A complex algebra \mathcal{A} is said to be self-adjoint if for every $f \in \mathcal{A}$, $\bar{f} \in \mathcal{A}$, where \bar{f} denotes the complex conjugate of f .

Theorem 7.25. Let X be a compact Hausdorff space and $\mathcal{A} \subseteq (X, \mathbb{C})$ a self-adjoint complex algebra which separates points and vanishes at no point of X . Then, \mathcal{A} is dense in $\mathcal{C}(X, \mathbb{C})$.

Proof. Let $f = u + iv \in \mathcal{A}$. Then $u = \frac{f + \bar{f}}{2} \in \mathcal{A}$. Let $x_1, x_2 \in X$ be distinct points. Then, due to Lemma 7.22, there is $f \in \mathcal{A}$ such that $f(x_1) = 0$ and $f(x_2) = 1$, then, for $u = \frac{f + \bar{f}}{2} \in \mathcal{A}$, $u(x_1) = 0$ and $u(x_2) = 1$. Finally, let $x_0 \in X$, then there is $f \in \mathcal{A}$ such that $f(x_0) \neq 0$, then, the function $g = \frac{\bar{f}(x_0)}{f(x_0)} f(x) \in \mathcal{A}$ is such that $g(x_0) > 0$, and thus $\Re(g)(x_0) > 0$.

If we denote $\mathcal{A}_R = \{\Re(f) \mid f \in \mathcal{A}\}$, then \mathcal{A}_R forms a real algebra that separates points and vanishes at no point of X . Therefore, for any $f = u + iv \in \mathcal{C}(X, \mathbb{C})$, there is a sequence of functions u_n, v_n converging uniformly to u and v respectively. As a result, $f_n = u_n + iv_n$ converges uniformly to f . This completes the proof. ■

Theorem 7.26. *Let X be a compact Hausdorff space. Then X is metrizable if and only if $\mathcal{C}(X, \mathbb{R})$ is separable.*

Proof.



Chapter 8

Baire Spaces and Dimension Theory

Definition 8.1 (Baire Space). A space X is said to be a *Baire space* if the following condition holds:

Given any countable collection $\{A_n\}$ of closed sets in X , each of which has empty interior in X , their union $\bigcup_{n=1}^{\infty} A_n$ also has empty interior in X .

Lemma 8.2. Let X be a topological space. Then X is Baire if and only if given any countable collection $\{U_n\}$ of open dense sets in X , their intersection $\bigcap_{n=1}^{\infty} U_n$ is dense in X .

Lemma 8.3. Let X be a complete metric space and $A_1 \supseteq A_2 \supseteq \cdots$ be a descending chain of nonempty closed sets in X with $\lim_{n \rightarrow \infty} \text{diam } A_n = 0$. Then, $\bigcap_{n=1}^{\infty} A_n$ is nonempty.

Proof. Using the Axiom of Choice, pick $x_n \in A_n$. It is not hard to show that these form a Cauchy sequence. Thus, must converge to some $x \in X$. Using simple arguments, it is easy to see that $x \in A_n$ for all $n \in \mathbb{N}$. Thus, $x \in \bigcap_{n=1}^{\infty} A_n$, and thus the intersection is nonempty. ■

Theorem 8.4. Let X be a topological space. Then, X is Baire if

- (a) X is compact and Hausdorff.
- (b) X is a complete metric space.

Proof. Let $\{U_n\}$ be a countable collection of open, dense subsets of X . Let $G = \bigcap U_n$. Let $x \notin G$ and U a neighborhood of x .

- (a) The intersection $U_1 \cap U$ is a nonempty open set in X , because U_1 is dense in X . Pick some point $x_1 \in U_1 \cap U$. Using the regularity of X , there is an open set V_1 containing x such that $\overline{V_1} \subseteq U_1$. Now, V_1 is an open set containing x_1 , consequently, has nonempty intersection with U_2 . As a result, there is a point $x_2 \in V_1 \cap U_2$ and proceeding similarly, we obtain a descending chain $\overline{V_1} \supseteq \overline{V_2} \supseteq \cdots$ of closed sets in X , all of which are contained in U . As a result, $V = \bigcap \overline{V_n}$ is contained in U and due to Cantor's

Intersection Theorem, is nonempty. Hence,

$$\emptyset \neq \bigcap_{n=1}^{\infty} \overline{V_n} \subseteq U \cap \bigcap_{n=1}^{\infty} U_n = U \cap G$$

thus G is dense in X .

- (b) When X is a complete metric space, we use a similar strategy as above. Just, instead of using regularity, we use the fact that any open ball contains a closed ball of arbitrarily small radius. Finally, we would have a descending sequence of nonempty closed sets $A_1 \supseteq A_2 \supseteq \cdots$ with $\lim_{n \rightarrow \infty} \text{diam } A_n = 0$, and due to the preceding lemma, their intersection is nonempty. ■

Lemma 8.5. *Any open subspace Y of a Baire space X is itself a Baire space.*

Proof. Let $\{A_n\}$ be a countable collection of closed sets with empty interiors in Y . Let $\overline{A_n}$ denote the closure of A_n in X . We contest that $\overline{A_n}$ has empty interior, for if $U \subseteq \overline{A_n}$, then it is not hard to show that $U \cap A_n \neq \emptyset$, from which it follows that $U \cap Y \subseteq A_n$, contradicting the fact that A_n has an empty interior.

Thus, $\bigcup \overline{A_n}$ has empty interior in X . Now, if $U \subseteq \bigcup A_n$ in Y , then $U \subseteq \bigcup \overline{A_n}$, and U is open in X since Y is open in X , a contradiction. Thus Y is Baire. ■

Corollary 8.6. *A locally compact Hausdorff space is Baire.*

Proof. Let X be a locally compact Hausdorff space and Y its one point compactification. Since Y is compact Hausdorff, it is Baire. Further, since X is an open subset of Y , it must be Baire. ■